

1 **Fire intensity, fire severity and burn severity: A brief review and**  
2 **suggested usage**

3

4

5 *Jon E. Keeley*<sup>A,B</sup>

6

7 <sup>A</sup> *USGS Western Ecological Research Center, Sequoia - Kings Canyon Field Station, 47050*

8 *Generals Highway, Three Rivers, CA 93271, USA*

9 <sup>B</sup> *Department of Ecology and Evolutionary Biology, University of California, Los Angeles,*

10 *CA 90095, USA*

11

12

13

14

15

16 E-mail: *jon\_keeley@usgs.gov*

17

18 *TOC: Contrary to some recent suggestions, fire intensity, fire severity, and burn severity are*

19 *terms that should be retained, but defined operationally; severity metrics may create confusion*

20 *when used to represent both fire/burn severity and ecosystem responses.*

21

22 **Abstract.** Several recent papers have suggested replacing the terminology of *fire intensity* and  
23 *fire severity*. Part of the problem with *fire intensity* is that it is sometimes used incorrectly to  
24 describe fire effects, when in fact it is justifiably restricted to measures of energy output.  
25 Increasingly the term has created confusion because some authors have restricted its usage to a  
26 single measure of energy output referred to as fireline intensity. This metric is most useful in  
27 understanding fire behavior in forests, but is too narrow to fully capture the multitude of ways  
28 fire energy affects ecosystems. Fire intensity represents the energy released during various  
29 phases of a fire and different metrics such as reaction intensity, fireline intensity, temperature,  
30 heating duration, and radiant energy are useful for different purposes. *Fire severity*, and the  
31 related term *burn severity*, has created considerable confusion because of recent changes in their  
32 usage. Some authors have justified this by contending that fire severity is defined broadly as  
33 ecosystem impacts from fire and thus is open to individual interpretation. I argue that based on a  
34 long history of empirical studies, fire severity is operationally defined as the loss of or change in  
35 organic matter aboveground and belowground, although the precise metric varies with  
36 management needs. Confusion arises because fire or burn severity is sometimes defined to also  
37 include ecosystem responses. *Ecosystem responses* include soil erosion, vegetation regeneration,  
38 restoration of community structure, faunal recolonisation, and a plethora of related response  
39 variables. Although some ecosystem responses are correlated with measures of fire or burn  
40 severity, many important ecosystem processes have either not been demonstrated to be predicted  
41 by severity indices or have been shown in some vegetation types to be unrelated to severity. This  
42 is a critical issue because fire or burn severity are readily measurable parameters, yet ecosystem  
43 responses are ultimately what are of most interest to resource managers.

44 **Additional keywords:** BAER, dNBR Landsat Thematic Mapper, Soil burn severity

## 45 **Introduction**

46 In recent papers dealing with postfire studies there has been a disturbing number that have  
47 acknowledged problems in terminology associated with fire intensity and fire severity (e.g.,  
48 Simard 1991; Parsons 2003; Jain et al. 2004; Lentile et al. 2006). These problems are perceived  
49 to be sufficiently problematical that alternative terminology has been proposed. Jain et al. (2004)  
50 suggested that these categories might best be replaced with a continuum of postfire changes,  
51 along the lines of Simard's (1991) space/time continuum of fire issues. It has also recently been  
52 suggested that fire intensity and severity be replaced with new categories such as "active fire  
53 characteristics" and "post-fire effects" (Lentile et al. 2006).

54 The present paper is prompted because of strong agreement about the problems in this  
55 terminology, but here I argue for retention of the original terminology as a valuable  
56 organizational tool. I believe that much of the confusion can be alleviated by clarification of the  
57 original operational definition of these terms and suggest a model that may help clarify the  
58 phenomena under consideration (Fig. 1). The emergence of remote imaging technology and its  
59 application to fire issues has contributed to some of the problems, in part because the speed of  
60 technology development has not always been in sync with our ability to relate it to useful  
61 purposes. It is argued that the basis of some of the problems has been the more recent  
62 introduction of the term burn severity and the extension of this term to include not just fire  
63 severity but what are here termed *ecosystem responses* (Fig. 1). These problems are illustrated  
64 with an example of the relationship of remote imaging signals to fire severity and ecosystem  
65 responses in southern California shrublands. The overriding goal is to point out those aspects of  
66 each term for which there has been general agreement and note those areas where further  
67 research and discussion are needed.

## 68 **Fire Intensity**

69 Fire intensity describes the physical process of fire releasing energy from organic matter. Thus, it  
70 would be logical to consider the usage of the term “intensity” in the field of physics, where it is  
71 defined as a measure of the time-averaged energy flux or in other words the energy per unit  
72 volume multiplied by the velocity at which the energy is moving; the resulting vector has the  
73 units of watt/m<sup>2</sup>. Rothermel’s (1972) reaction intensity, which represents the heat source in his  
74 firespread model, is consistent with this definition. However, fire science is like many other  
75 fields that have demonstrated other needs for the term “intensity.”

76 One example is fireline intensity, which is the rate of heat transfer per unit length of the fire  
77 line (kW m<sup>-1</sup>) (Byram 1959). This represents the radiant energy release in the flaming front and  
78 is an important characteristic for propagation of a fire and thus is critical information for fire  
79 suppression activities and has been incorporated into fire danger rating calculations (Salazar and  
80 Bradshaw 1986; Hirsch and Martell 1996; Weber 2001). Increasingly, fireline intensity is  
81 presented in the literature as the only appropriate measure for fire intensity (e.g., Johnson 1992;  
82 Michaletz and Johnson 2003; Chatto and Tolhurst 2004; Sugihara et al. 2006), but this is  
83 misleading because it fails to acknowledge that for many fire scientists other measures of energy  
84 release from fires provide more useful metrics.

85 Fireline intensity is most frequently used in forested ecosystems as there is a well developed  
86 literature showing a relationship between fireline intensity or flame length and scorching height  
87 of conifer crowns and other biological impacts of fire. However, some fire effects are more  
88 closely tied to other fire intensity metrics. For example, modeling duff consumption requires  
89 understanding smoldering combustion., which is more related to temperatures at the soil surface  
90 and the duration of heating than to fireline intensity (Ryan and Frandsen 1991; Hartford and

91 Frandsen 1992; Valette et al. 1994; Miyanishi 2001). Even with tree mortality, fireline intensity  
92 often can not explain mortality patterns since mortality may be more a function of total heat  
93 output reflected in flame residence time or a function of smoldering combustion in the duff after  
94 the flame front passes (Wade 1993; Sackett et al. 1996). Also, the development of non-wettable  
95 layers in soil may be more closely related to duration of soil heating (DeBano 2000), and  
96 survival of seed banks or rhizomes may be closely tied to duration of heating as well as  
97 maximum soil temperatures (Beadle 1940; Flinn and Wein 1977; Auld and O'Connell 1991;  
98 Bradstock and Auld 1995; Brooks 2002). Measurements of these other metrics are often  
99 required since fireline intensity may be weakly correlated with maximum temperature or heating  
100 duration (Bradstock and Auld 1995; Keeley and McGinnis 2007). This should be no surprise  
101 since very little radiant or convected heat from combustion of aerial fuels may be transferred to  
102 the soil, and often soil temperatures are more dependent on consumption of fine fuels on the  
103 surface (Bradstock and Auld 1995). Although fireline intensity provides information for fire  
104 managers involved in fire containment, temperature and duration of heating (residence time) may  
105 be far more critical information for managers concerned with prescribed burning conditions  
106 required to retain sensitive ecosystem components. In addition, the future for fire science will be  
107 heavily influenced by remote imaging technologies and these may not always scale with fireline  
108 intensity (Smith et al. 2005). Other metrics, such as radiative energy appear to be a more readily  
109 measurable metric for fire intensity in remote imaging studies of fire impacts (Wooster et al.  
110 2003; Dennison et al. 2006).

111 Another reason for not discounting other metrics of fire intensity is that fireline intensity has  
112 important limitations, particularly in how it is measured and its ability to make cross ecosystem  
113 comparisons. Byram's fireline intensity assumes that available fuel weight reflects fuels entirely

114 consumed during the flaming phase of combustion as the flame front passes. This metric  
115 excludes glowing combustion or post-frontal smoldering, which may continue for many hours  
116 after the front passes. Thus, fireline intensity requires that one distinguish fuels consumed by the  
117 flaming front from the total fuel consumption. However, fuel consumption usually is estimated  
118 as the difference between pre-and post fire fuel inventories, and this inflates estimates of fireline  
119 intensity (Alexander 1982; Scott and Reinhardt 2001). Because of these difficulties the majority  
120 of papers reporting fireline intensity do not measure it directly, rather they utilize surrogate  
121 measures that are assumed to be allometrically related. Typically, flame length is used and much  
122 work has gone into methodology development for making such measurements (Ryan 1981;  
123 Finney and Martin 1992). Empirical studies show there is a significant relationship between  
124 flame length and fireline intensity in forest and shrubland ecosystems (Andrews and Rothermel  
125 1982; Johnson 1992; Wade 1993; Burrows 1995; Fernandes et al. 2000). However, in vegetation  
126 with a mixture of fine fuels and woody fuels such as palmetto understories or grasslands and  
127 savanna forests the relationship is not always reliable (Nelson and Adkins 1986; Catchpole et al.  
128 1993; Keeley and McGinnis 2007). Cheney (1990) found that fireline intensity is system  
129 dependent and fires of identical intensities in different fuel beds will have very different flame  
130 lengths. Thus, flame length is only applicable to fuel types with the same fuel structure  
131 characteristics.

132 *In summary, fire intensity represents the energy released during various phases of the fire*  
133 *and no single metric captures all of the relevant aspects of fire behavior. Different metrics,*  
134 *including reaction intensity, fireline intensity, temperature, residence time, radiant energy and*  
135 *others are useful for different purposes.*

136

## 137 **Fire Severity**

138 The term fire severity was born out of the need to provide a description of how fire intensity  
139 affected ecosystems, particularly following wildfires where direct information on fire intensity  
140 was absent. Some definitions of fire severity have been rather general statements about broad  
141 impacts of fires, e.g., the degree of environmental change caused by fire (e.g., White and Pickett  
142 1985; Simard 1991; Jain et al. 2004, NWCG 2006), and consequently have not lent themselves to  
143 operationally useful metrics. However, most empirical studies that have attempted to measure  
144 fire severity have had a common basis that centers on the loss or decomposition of organic  
145 matter, both aboveground and belowground. Aboveground metrics such as crown volume scorch  
146 used in forests or twig diameter remaining on terminal branches used in forests and shrublands  
147 are indicators of biomass loss (e.g., van Wagner 1973; Moreno and Oechel 1989; Tolhurst 1995;  
148 Dickinson and Johnson 2001). Soil characteristics include the loss of the litter and duff layers  
149 and ash characteristics, all of which reflect the level of organic matter consumed (Wells et al.  
150 1979; Stronach and McNaughton 1989; Neary et al. 1999; Ice et al. 2004).

151 One of the first metrics for fire severity that captured the essence of how it subsequently has  
152 been used empirically was that proposed by Ryan and Noste (1985). They maintained that any  
153 metric for fire severity needed to consider the immediate impacts of heat pulses aboveground and  
154 belowground, which they noted were directly related to fire intensity. They developed an index  
155 that comprised a matrix of vegetation and soil impacts reflecting the degree of organic matter  
156 consumed, which in most studies has been simplified to categories of fire severity (Table 1).  
157 They, and others (e.g., Cram et al. 2006), have found this index does capture the fire intensity  
158 signal, and appears to be a function of fireline intensity, residence time (heating duration) and  
159 soil and plant dryness (Chatto and Tolhurst 2004). Of course other factors such as prefire species

160 composition, stand age, topography, substrate, and climate will all have some effect on how fire  
161 intensity translates into fire severity.

162 Many studies that report fire severity have utilized an index similar to Table 1 or at least an  
163 index based on the concept of organic matter loss, such as crown volume scorch, and these have  
164 been shown to be correlated with measures of fire intensity (Buckley 1993; Williams et al. 1998;  
165 Catchpole 2000). Depending on the focus of the study they may report only on vegetation or on  
166 soils. For example, the BAER (Burned Area Emergency Response (formerly Rehabilitation)  
167 assessment, which is conducted by U.S. federal government agencies has traditionally focused  
168 on soil changes induced by fire and has often referred to this as the *soil burn severity* assessment  
169 (see Burn severity section). In these soil assessments the metric is largely based on loss of soil  
170 organic matter or deposition of ash from the aboveground combustion of biomass (Lewis et al.  
171 2006). Other parameters that are sometimes included in this assessment of fire severity impacts  
172 to soils include changes in soil structure, increased hydrophobicity, and iron oxidation, many of  
173 which are indirectly tied to organic matter decomposition as well. Of course the purpose of such  
174 assessments is not because of any perceived need to determine organic matter loss, but rather  
175 because it is presumed that these are keys to other impacts (discussed under Ecosystem  
176 response). Whether or not studies have used the Ryan and Noste (1985) index in its entirety,  
177 most have used metrics that depend on loss of organic matter and in that respect share the same  
178 functionality as that index.

179 Remote imaging studies have found a good correlation between the LANDSAT signals,  
180 particularly the Normalised Difference Vegetation Index (NDVI), and fire severity estimates  
181 based on biomass loss (e.g., Turner et al. 1994; Miller and Yool 2002; Conard et al. 2002; Chafer  
182 et al. 2004). Much of this work has been done in forests and woodlands and studies that have

183 sampled more broadly have found that the vegetation type markedly influences the detection of  
184 fire severity (Hammill and Bradstock 2006).

185 Plant mortality, which is also a measure of biomass loss, is often included in fire severity  
186 metrics, or sometimes the fire severity metric is based entirely on mortality (e.g., Chappell and  
187 Agee 1996; Larson and Franklin 2005). Numerous studies have shown that fire intensity is  
188 correlated with mortality and other measures of biomass loss such as crown scorch (e.g., Wade  
189 1993; McCaw et al. 1997). Tree mortality has been widely used in conifer forests in North  
190 America that historically have been exposed to low severity or mixed severity fire regimes where  
191 there is substantial tree survivorship. In these forests the dominant trees are non-sprouting  
192 species so that aboveground mortality reflects mortality of the entire genet. One limitation to  
193 using mortality is that it sometimes is not evident for a year or more after a fire event. Where the  
194 use of this metric becomes very problematical is when it is applied to understory species in many  
195 forest types or to dominant species in crown-fire ecosystems such as shrublands. In these species  
196 the aboveground ramets are nearly always killed, but some percentage survive belowground. A  
197 problem is created when the degree of resprouting is incorporated into the mortality index  
198 because resprouting is often not related to fire intensity. Many species are innately incapable of  
199 resprouting (Keeley 1981) and within resprouting species there is substantial variation in  
200 resprouting capacity that is related to species-specific differences (Vesk and Westoby 2004) and  
201 plant age (Keeley 2006a). Without considering site to site variation in prefire species  
202 composition, resprouting should not be included as a measure of fire severity and as discussed  
203 below, is best viewed as an ecosystem response variable.

204 *In summary, fire severity refers to loss or decomposition of organic matter aboveground and*  
205 *belowground. Metrics for this parameter vary with the ecosystem. Including mortality is*

206 *consistent with the definition of fire severity as a loss of organic matter, however, it is only*  
207 *advisable when dealing with forest trees that lack any resprouting capacity. Fire severity is*  
208 *correlated with fire intensity.*

## 209 **Burn severity**

210 The term *burn severity* has gained popularity in recent years but it has caused some confusion  
211 because it is often used interchangeably with fire severity, and often using metrics consistent  
212 with fire severity measurement (e.g., White et al. 1996; Turner et al. 1999; Rogan and Franklin  
213 2001). In the U.S. BAER (Burned Area Emergency Response) assessments, the term burn  
214 severity has largely replaced fire severity although the metric is very similar and is largely based  
215 on loss of organic matter in the soil and aboveground organic matter conversion to ash. In the  
216 recent “Glossary of Wildland Fire Terminology” the term burn severity is restricted to the loss of  
217 organic matter in or on the soil surface (NWCG 2006), and in this respect represents what  
218 BAER assessments term “soil burn severity” (Parsons 2003).

219 Remote sensing applications to assessing burned areas typically use the term burn severity  
220 rather than fire severity, and as remote sensing has increased in burned area assessments, so has  
221 the use of the term burn severity. In some of the initial studies of remote sensing applications to  
222 burned area assessments the term burn severity was used for the index calculated from the  
223 satellite sensors (van Wagtenonk et al. 2004). Various sensors (e.g., MODIS, AVIRIS) have  
224 been tested for their ability to match field measurements of severity and the Landsat Thematic  
225 Mapper sensor is widely accepted as most appropriate for this task (van Wagtenonk et al. 2004;  
226 Epting et al. 2005; Brewer et al. 2005; Cocks et al. 2005; Chuvieco et al. 2006; but c.f. Roy et al.  
227 2006; Kokaly et al. 2007). These remote sensing data are used to generate an index known as

228 the differenced Normalized Burn Ratio (dNBR), which is a preferable term over burn severity as  
229 it keeps separate the remote imaging index from surface measurements of the burned site.

230 BAER assessments are now commonly expedited by the use of satellite sensing data that use  
231 the dNBR index to produce a burn severity map of conditions on the ground, and this is termed  
232 the Burned Area Reflectance Classification (BARC). There appears to be a reasonably good  
233 correlation between these BARC map categories and field assessments of fire severity (Bobbe et  
234 al. 2004; Robichaud et al. 2007b), however, since the assessments must be done very soon after  
235 the fire it is not always possible to coordinate satellite pass-over with clear skies.

236 In many remote sensing studies field validation of the method has utilized metrics of fire  
237 severity, i.e., organic matter loss through combustion or mortality viz a viz Ryan and Noste  
238 (1985), although sometimes using the term burn severity (White et al. 1996; Rogan and Franklin  
239 2001; Miller and Yool 2002; Chafer et al. 2004; Hammill and Bradstock 2006; Roldán-Zamarrón  
240 et al. 2006).

241 In recent studies utilizing remote sensing indices, field validation has used the term burn  
242 severity in a way that diverges from the concept of fire severity as a measure of just organic  
243 matter loss, rather in these studies burn severity defines a much broader collection of attributes  
244 that include both fire severity and ecosystem responses (van Wagtendonk et al. 2004; Epting et  
245 al. 2005; Cocke et al. 2005; Chuvieco et al. 2006). This approach is described as the *composite*  
246 *burn index* and it is designed to provide a single index that represents many different phenomena  
247 of interest to land managers (Key and Benson 2006). The composite index combines fire  
248 severity metrics and ecosystem recovery that includes resprouting of herbs, shrubs and hardwood  
249 trees, and seedling colonization. Recent studies of several major fires in southern California raise  
250 concerns about the value of combining fire severity and ecosystem responses into a single

251 “composite” index (Box 1). These studies show that while dNBR is significantly correlated with  
252 field measurements of fire severity, this signal is not necessarily a good predictor of ecosystem  
253 responses. This is critical because the remote imaging signal is most important to land managers  
254 only as far as it is a predictor of ecosystem responses. The potential for remote sensing  
255 techniques to contribute to postfire management has not yet been fully realized and it is  
256 suggested that this will develop best if we parse out the separate contributions of fire severity and  
257 ecosystem response (Fig. 1).

258 *In summary, when the term burn severity is used interchangeably with fire severity it may*  
259 *lead to some minor confusion but is not a significant problem. However, where the term has been*  
260 *defined to include fire severity and ecosystem responses it may lead to a significant amount of*  
261 *confusion as it has the potential for confounding factors with different effects. It is recommended*  
262 *that fire severity and ecosystem responses be evaluated separately.*

## 263 **Ecosystem Response**

264 Fire intensity, fire severity and burn severity are operationally tractable measures, but they are  
265 largely of value only so far as they can predict ecosystem responses such as soil erosion or  
266 natural revegetation. In addressing this issue, fire scientists may take one of two approaches: the  
267 descriptive approach or the process-based approach (Johnson and Miyanishi 2001; Michaletz and  
268 Johnson 2003). The former yields statistical descriptions of relationships between for example  
269 fire intensity and fire severity, or fire severity and ecosystem responses, and this is often the only  
270 approach available when studying impacts of wildfires. Under more controlled experimental  
271 conditions one can use the process-based approach that studies the direct path from measures of  
272 fire intensity to fire severity or from fire intensity to ecosystem response variables and tests  
273 underlying mechanisms. Regardless of the path studied, it is clear that many biotic and abiotic

274 factors also enter into the relationship between fire intensity and ecosystem response (e.g.,  
275 Peterson and Ryan 1986; Neary et al. 1999; Moody and Martin 2001; Pérez-Cabello et al. 2006).

276 Statistical studies show correlations between fire intensity and fire severity metrics (e.g.,  
277 McCaw et al. 1997) and between different measures of fire severity and ecosystem responses.  
278 For example, in forests it has been shown that fire severity is tied to forest recovery and alien  
279 plant invasion (Turner et al. 1999; Wang and Kembell 2003) and belowground changes in fauna  
280 and flora (Neary et al. 1999). In crown-fire forests and shrublands, increased fire severity has  
281 been correlated with decreased resprouting of herbs and shrubs (Flinn and Wein 1977; Keeley  
282 2006). Fire severity has also been correlated with ecosystem responses such as species richness  
283 and patterns of seedling recruitment (Whelan 1995; Bond and van Wilgen 1996; Ryan 2002;  
284 Keeley et al. 2005; Johnstone and Chapin 2006). In some shrublands, high fire severity is  
285 correlated with reduced alien plant invasion (Keeley 2006). In Canadian boreal forests fire  
286 severity may be correlated with long lasting impacts on forest regeneration and carbon storage  
287 (Lecomte et al. 2006). On the other hand in some ecosystems important responses such as  
288 vegetative regeneration or resprouting after fire are not correlated with fire severity measures on  
289 the ground or remote sensing indices (Box 1).

290 Process-based studies can provide a mechanistic basis for translating fire intensity measures  
291 directly into fire severity impacts such as tree mortality as well as ecosystem responses such as  
292 erosion. One of the clearest examples is the use of heat transfer models of the flame and plume  
293 heat into a plant to account for tree mortality patterns (Gill and Ashton 1968; Dickinson and  
294 Johnson 2001). Mercer et al. (1994) demonstrated that seed survival in woody fruits was  
295 predicted by a mathematical model that used heat-flow equations with time-dependent  
296 temperature inputs and used this model to predict seed survival in the field. Temperature

297 response curves for seed survival, when coupled with field measures of fire intensity, also  
298 provide predictive models for subsequent seedling recruitment (Keeley and McGinnis 2007).

299 A major reason for postfire assessments of fire or burn severity is because it is believed to be  
300 an important indicator of the potential for water runoff and erosion (Robichaud et al. 2000;  
301 Wilson et al. 2001; Ruiz-Gallardo et al. 2004; Lewis et al. 2006). Indeed, it is sometimes stated  
302 that these severity measurements are indicators of changes in soil hydrologic function (Parsons  
303 2003; Ice et al. 2004). Conceptually this inference is logical based on various types of indirect  
304 evidence. For example, loss of aboveground biomass exposes more soil surface, which increases  
305 the kinetic force of precipitation on the soil surface and that can increase overland flow (Moody  
306 and Martin 2001). Also, loss of soil organic matter alters the binding capacity of soil and results  
307 in other structural changes that can affect erosional processes (Hubbert et al. 2006). Postfire  
308 increases in soil water repellency due to hydrophobic soil layers is tied, albeit sometimes weakly,  
309 to fire severity (Robichaud 2000; Lewis et al. 2006), although in some ecosystems soil  
310 hydrophobicity is unrelated to fire severity (Cannon et al. 2001; Doerr et al. 2006).

311 In general, there is little direct evidence that fire severity measurements are a reliable  
312 indicator of specific changes in hydrologic or other ecosystem functions (Robichaud et al. 2000;  
313 Gonzalez-Pelayo et al. 2006), and some even suggest that fire severity classifications are  
314 unsuitable for predicting fire impacts on soil hydrological responses (Doerr et al. 2006). The  
315 primary reason is that ecological responses such as erosion, overland water flow and debris flows  
316 are affected as much by topography, soil type, rates of weathering, fire-free interval, and  
317 precipitation as they are by fire severity (Moody and Martin 2001; Cannon et al. 2001; Nearing  
318 et al. 2005). In short, the factors responsible for hydrologic responses to fire are multi-factorial  
319 and until we have better mechanistic models explaining these phenomena it would be prudent to

320 keep separate the metric for fire or burn severity from inferred ecosystem responses. Applied  
321 efforts focused on this include Erosion Risk Management Tool (ERMiT) (Robichaud et al.  
322 2007a).

323 *Ecosystem responses include those processes that are differentially affected by fire intensity,*  
324 *measured either directly, or indirectly with fire severity metrics, and include erosion, vegetation*  
325 *regeneration, faunal recolonisation, restoration of community structure and a plethora of other*  
326 *response variables. Predicting how fire intensity or severity will affect these responses is critical*  
327 *to postfire management.*

## 328 **Conclusions**

329 A summary of the appropriate and inappropriate use of these terms is in Table 2. Fire intensity is  
330 the energy output from fire and should not be used to describe fire effects. Fire severity and burn  
331 severity have been used interchangeably and operationally have generally emphasized degrees of  
332 organic matter loss or decomposition both aboveground and belowground. Both are positively  
333 correlated with fire intensity. Significant confusion has arisen from rather broad definitions for  
334 fire or burn severity that include ecosystem responses. Another source of confusion has arisen by  
335 using these terms for remote sensing indices and separate terms such as BARC or dNBR are  
336 preferable. Ecosystem responses include vegetative regeneration and faunal recolonization as  
337 well as abiotic watershed hydrologic processes. Some of these have been directly correlated with  
338 fire intensity and others indirectly with fire or burn severity metrics. Ecosystem responses may  
339 be positive, negative or neutral in their response to fire intensity and severity.

340 This approach has value for resource managers because it emphasizes the distinction between  
341 measures of severity after a fire and the resource impact of the fire. Most managers are not  
342 specifically interested in severity measures per se, but rather the extent to which they reflect

343 potential ecosystem responses. Metrics that combine burn severity and measures of vegetative  
344 recovery can provide misinformation when those measures are not correlated. It is recommended  
345 that field measurements of severity be restricted to measures of organic matter loss, such as  
346 canopy scorch or ash deposition, and these be analyzed separately from measures of ecosystem  
347 response such as vegetative regeneration. Mortality needs to be evaluated with consideration of  
348 species-specific traits. Mortality is a straightforward measure in most conifer dominated forests  
349 but in other ecosystems it can only be evaluated in the context of prefire community composition  
350 because of species-specific differences in resprouting capacity.

351

352

353

354

### 355 **Acknowledgments**

356 This manuscript has greatly benefited from discussion with, and/or comments on an earlier draft  
357 by, the following colleagues: Jan Beyers, James Grace, Carl Key, Jay Miller, Jason Moghaddas,  
358 Annette Parsons, David L. Peterson, Karen Phillips, Bill Romme, Kevin Ryan, Hugh Safford,  
359 Phillip van Mantgem and Marti Witter Thanks to Jeff Eidenshink for providing remote sensing  
360 dNBR indices. This research was made possible through funding of the Joint Fire Science  
361 Program Project 04-1-2-01. Any use of trade, product, or firm names in this publication is for  
362 descriptive purposes only and does not imply endorsement by the U.S. government.

363

364

365

366

367 **References**

- 368 Alexander ME (1982) Calculating and interpreting forest fire intensities. *Canadian Journal of*  
369 *Botany* **60**, 349-357.
- 370 Andrews PL, Rothermel RC (1982) 'Charts for interpreting wildland fire behavior  
371 characteristics.' USDA Forest Service, Intermountain Forest and Range Experiment Station,  
372 General Technical Report INT-131. (Ogden, UT).
- 373 Auld TD, O'Connell MA (1991) Predicting patterns of post-fire germination in 25 eastern  
374 Australian Fabaceae. *Australian Journal of Ecology* **16**, 53-70.
- 375 Beadle NCW (1940) Soil temperatures during forest fires and their effect on the survival of  
376 vegetation. *Journal of Ecology* **28**, 180-192.
- 377 Bobbe T, Finco MV, Quayle B, Lannom K, Sohlberg R, Parsons A (2004) 'Field measurements  
378 for the training and validation of burn severity maps from spaceborne, remotely sensed  
379 imagery.' Joint Fire Science Program (Boise, ID) ([http://jfsp.nifc.gov/news/doc/highlight5-](http://jfsp.nifc.gov/news/doc/highlight5-04.pdf)  
380 [04.pdf](http://jfsp.nifc.gov/news/doc/highlight5-04.pdf))
- 381 Bond WJ, van Wilgen BW (1996) 'Fire and plants.' (Chapman & Hall: New York, NY)
- 382 Bradstock RA, Auld TD (1995) Soil temperatures during experimental bushfires in relation to  
383 fire intensity: consequences for legume germination and fire management in south-eastern  
384 Australia. *Journal of Applied Ecology* **32**, 76-84.
- 385 Brewer KC, Winne JC, Redmond RL, Opitz DW, Mangrich MV (2005) Classifying and  
386 mapping wildfire severity: A comparison of methods. *Photogrammetric Engineering and*  
387 *Remote Sensing* **7**, 1311-1320.
- 388 Brooks ML (2002) Peak fire temperatures and effects on annual plants in the Mojave Desert.  
389 *Ecological Applications* **12**, 1088-1102.

- 390 Buckley AJ (1993) 'Fuel reducing regrowth forests with a wiregrass fuel type: fire behaviour  
391 guide and prescriptions.' Department of Conservation and Environment (Melbourne,  
392 Victoria, Australia)
- 393 Burrows ND (1995) A framework for assessing acute impacts of fire in jarrah forests for  
394 ecological studies. *CALM Science Supplement* **4**, 59-66.
- 395 Byrum GM (1959) Combustion of forest fuels. In 'Forest fire: Control and use.' (Ed KP Davis)  
396 pp. 61-89. (McGraw-Hill: New York, NY)
- 397 Cannon SH, Kirkham RM, Parise M (2001) Wildfire-related debris-flow initiation processes,  
398 Storm King Mountain, Colorado. *Geomorphology* **39**, 171-188.
- 399 Catchpole EA, Catchpole WR, Rothermel RC (1993) Fire behavior experiments in mixed fuel  
400 complexes. *International Journal of Wildland Fire* **3**, 45-57.
- 401 Catchpole W (2000) The international scene and its impact on Australia. In 'Fire! The Australian  
402 experience' pp. 137-148. Australian Academy of Technological Sciences and Engineering  
403 Limited (Canberra, ACT, Australia)
- 404 Chafer CJ, Noonan M, Mcnaught E (2004) The post-fire measurement of fire severity and  
405 intensity in the Christmas 2001 Sydney wildfires. *International Journal of Wildland Fire* **13**,  
406 227-240.
- 407 Chappell CB, Agee JK (1996) Fire severity and tree seedling establishment in *Abies magnifica*  
408 forests, southern Cascades, Oregon. *Ecological Applications* **6**, 628-640.
- 409 Chatto K, Tolhurst KG (2004) 'A review of the relationship between fireline intensity and the  
410 ecological and economic effects of fire, and methods currently used to collect fire data.' Fire  
411 Management, Department of Sustainability and Environment (Victoria, Australia)
- 412 Cheney P (1990) Quantifying bushfires. *Mathematical Computer Modelling* **13**, 9-15.

- 413 Chuvieco E, Riano D, Danson FM, Martin P (2006) Use of a radiative transfer model to simulate  
414 the postfire spectral response to burn severity. *Journal of Geophysical Research* **111**,  
415 G04S09:1-15.
- 416 Cocks AE, Fule PZ, Crouse JE (2005) Comparison of burn severity assessments using  
417 Differenced Normalized Burn Ratio and ground data. *International Journal of Wildland Fire*  
418 **14**, 189-198.
- 419 Conard SG, Sukhinin AI, Stocks BJ, Cahoon DR, Davidenko EP, Ivanova GA (2002)  
420 Determining effects of area burned and fire severity on carbon cycling and emissions in  
421 Siberia. *Climatic Change* **55**, 197-211.
- 422 Cram DS, Baker TT, Boren J (2006) 'Wildland fire effects in silviculturally treated vs. untreated  
423 stands of New Mexico and Arizona.' USDS Forest Service, Rocky Mountain Research  
424 Station, Research Paper RMRS-RP-55. (Fort Collins, CO)
- 425 DeBano LF (2000) Water repellency in soils: a historical overview. *Journal of Hydrology* **231-**  
426 **232**, 4-32.
- 427 Dennison PE, Charoensiri K, Roberts DA, Peterson SH, Green RO (2006) Wildfire temperature  
428 and land cover modeling using hyperspectral data. *Remote Sensing of Environment* **100**,  
429 212-222.
- 430 Dickinson MB, Johnson EA (2001) Fire effects on trees. In 'Forest fires: behavior and ecological  
431 effects'. (Eds EA Johnson and K Miyanishi) pp. 477-525. (Academic Press: San Francisco,  
432 CA)
- 433 Doerr SH, Shakesby RA, Blake WH, Chafer CJ, Humphreys GS, Wallbrink PJ (2006) Effects of  
434 differing wildfire severities on soil wettability and implications for hydrological response.  
435 *Journal of Hydrology* **319**, 295-311.

- 436 Epting J, Verbyla D, Sorbel B (2005) Evaluation of remotely sensed indices for assessing burn  
437 severity in interior Alaska using Landsat TM and ETM+. *Remote Sensing of Environment*  
438 **96**, 328-339.
- 439 Fernandes PM, Catchpole WR, Rego FC (2000) Shrubland fire behaviour modelling with  
440 microplot data. *Canadian Journal of Forest Research* **30**, 889-899.
- 441 Finney MA, Martin RE (1992) Calibration and field testing of passive flame height sensors.  
442 *International Journal of Wildland Fire* **2**, 115-122.
- 443 Flinn MA, Wein RW (1977) Depth of underground plant organs and theoretical survival during  
444 fire. *Canadian Journal of Botany* **55**, 2550-2554.
- 445 Gill AM, Ashton DH (1968) The role of bark type in relative tolerance to fire of three Central  
446 Victorian eucalypts. *Australian Journal of Botany* **16**, 491-498.
- 447 González-Pelayo O, Andreu V, Campo J, Gimeno-García E, Rubio JL (2006) Hydrological  
448 properties of a Mediterranean soil burned with different fire intensities. *Catena* **68**, 186-193.
- 449 Hammill KA, Bradstock RA (2006) Remote sensing of fire severity in the Blue Mountains:  
450 influence of vegetation type and inferring fire intensity. *International Journal of Wildland*  
451 *Fire* **15**, 213-226.
- 452 Hartford RA, Frandsen WH (1992) When it's hot, it's hot ... or maybe it's not! (surface flaming  
453 may not portend extensive soil heating). *International Journal of Wildland Fire* **2**, 139-144.
- 454 Hirsch KG, Martell DL (1995) A review of initial attack fire crew productivity and effectiveness.  
455 *International Journal of Wildland Fire* **6**, 199-215.
- 456 Hubbert KR, Preisler HK, Wohlgemuth PM, Graham RC, Narog MG (2006) Prescribed burning  
457 effects on soil physical properties and soil water repellency in a steep chaparral watershed,  
458 southern California, USA. *Geoderma* **130**, 284-298.

- 459 Ice GG, Neary DG, Adams PW (2004) Effects of wildfire on soils and watershed processes.  
460 *Journal of Forestry* **102**, 16-20.
- 461 Jain T, Pilliod D, Graham R (2004) Tongue-tied. *Wildfire* **4**, 22-36.
- 462 Johnson EA (1992) 'Fire and vegetation dynamics: studies from the North American boreal  
463 forest.' (Cambridge University Press: Cambridge, U.K.)
- 464 Johnson EA, Miyanishi K (2001) Strengthening fire ecology's roots. In 'Forest fires. Behavior  
465 and ecological effects'. (Eds EA Johnson and K Miyanishi) pp. 1-9. (Academic Press: San  
466 Diego, CA)
- 467 Johnstone JF, F.S. Chapin I (2006) Effects of soil burn severity on post-fire tree recruitment in  
468 boreal forest. *Ecosystems* **9**, 14-31.
- 469 Keeley JE (1981) Reproductive cycles and fire regimes. In 'Proceedings of the conference fire  
470 regimes and ecosystem properties'. (Eds HA Mooney, TM Bonnicksen, NL Christensen, JE  
471 Lotan and WA Reiners) pp. 231-277. USDA Forest Service, General Technical Report WO-  
472 3. (Washington, D.C.)
- 473 Keeley JE (2006a) Fire severity and plant age in postfire resprouting of woody plants in sage  
474 scrub and chaparral. *Madroño* **53**, 373-379.
- 475 Keeley JE (2006b) Fire management impacts on invasive plant species in the western United  
476 States. *Conservation Biology* **20**, 375-384.
- 477 Keeley, J.E., T. Brennan, and A.H. Pfaff. Fire severity and ecosystem responses following crown  
478 fires in California shrublands. *Ecological Applications*. *In press*.
- 479 Keeley JE, Fotheringham CJ, Baer-Keeley M (2005) Determinants of postfire recovery and  
480 succession in mediterranean-climate shrublands of California. *Ecological Applications* **15**,  
481 1515-1534.

- 482 Keeley JE, McGinnis T (2007) Impact of prescribed fire and other factors on cheatgrass  
483 persistence in a Sierra Nevada ponderosa pine forest. *International Journal of Wildland Fire*  
484 **16**, 96-106.
- 485 Key CH, Benson NC (2006) 'Landscape assessment: sampling and analysis methods.' USDA  
486 Forest Service, Rocky Mountain Research Station (Ogden, UT)
- 487 Kokaly RF, Rockwell BW, Haire SL, King TVV (2007) Characterization of post-fire surface  
488 cover, soils, and burn severity at the Cerro Grande Fire, New Mexico, using hyperspectral  
489 and multipsectral remote sensing. *Remote Sensing of Environment* **106**, 305-325.
- 490 Larson AJ, Franklin JF (2005) Patterns of conifer tree regeneration following an autumn wildfire  
491 event in the western Oregon Cascade Range, USA. *Forest Ecology and Management* **218**,  
492 25-36.
- 493 Lecomte N, Simard M, Fenton N, Bergeron Y (2006) Fire severity and long-term ecosystem  
494 biomass dynamics in coniferous boreal forests of eastern Canada. *Ecosystems* **9**, 1215-1230.
- 495 Lentile LB, Holden ZA, Smith AMS, Falkowski MJ, Hudak AT, Morgan P, Lewis SA, Gessler  
496 PE, Benson NC (2006) Remote sensing techniques to assess active fire characteristics and  
497 post-fire effects. *International Journal of Wildland Fire* **18**, 319-345.
- 498 Lewis SA, Wu JQ, Robichaud PR (2006) Assessing burn severity and comparing soil water  
499 repellency, Hayman Fire, Colorado. *Hydrological Processes* **20**, 1-16.
- 500 Ludwig JA, Bastin GN, Wallace JF, McVicar TR (2007) Assessing landscape health by scaling  
501 with remote sensing: when is it not enough? *Landscape Ecology* **22**, 163-169.
- 502 McCaw WL, Smith RH, Neal JE (1997) Prescribed burning of thinning slash in regrowth stands  
503 of karri (*Eucalyptus diversicolor*). 1. Fire characteristics, fuel consumption and tree damage.  
504 *International Journal of Wildland Fire* **7**, 29-40.

- 505 Mercer GN, Gill AM, Weber RO (1994) A time-dependent model of fire impact on seed survival  
506 in woody fruits. *Australian Journal of Botany* **42**, 71-81.
- 507 Michaletz ST, Johnson EA (2003) Fire and biological processes. *Journal of Vegetation Science*  
508 **14**, 622-623.
- 509 Miller JD, Yool SR (2002) Mapping forest post-fire canopy consumption in several overstory  
510 types using multi-temporal Landsat TM and ETM data. *Remote Sensing of Environment* **82**,  
511 481-496.
- 512 Miyanishi K (2001) Duff consumption. In 'Forest fires: behavior and ecological effects'. (Eds EA  
513 Johnson and K Miyanishi) pp. 437-475. (Academic Press: San Francisco, CA)
- 514 Moody JA, Martin PA (2001) Initial hydrologic and geomorphic response following a wildfire in  
515 the Colorado front range. *Earth Surface Processes and Landforms* **26**, 1049-1070.
- 516 Moreno JM, Oechel WC (1989) A simple method for estimating fire intensity after a burn in  
517 California chaparral. *Acta Oecologica* **10**, 57-68.
- 518 NWCG (2006) Glossary of wildland fire terminology. National Wildfire Coordinating Group,  
519 Incident Operations Standards Working Team (<http://www.nwec.gov>; accessed February  
520 2007)
- 521 Nearing MA, Jetten V, Baffaut C, Cerda O, Couturier A, Hernandez M, Le Bissonnais Y,  
522 Nichols MH, Nunes JP, Renschler CS, Souchère V, van Oost K (2005) Modeling response  
523 of soil erosion and runoff to changes in precipitation and cover. *Catena* **61**, 131-154.
- 524 Neary DG, Klopatek CC, DeBano LF, Ffolliott PF (1999) Fire effects on belowground  
525 sustainability: a review and synthesis. *Forest Ecology and Management* **122**, 51-71.
- 526 Nelson RM, Jr., Adkins CW (1986) Flame characteristics of wind-driven surface fires. *Canadian*  
527 *Journal of Forest Research* **16**, 1293-1300.

- 528 Parsons A (2003) 'Burned Area Emergency Rehabilitation (BAER) soil burn severity definitions  
529 and mapping guidelines. Draft.' USDA Forest Service, Rocky Mountain Research Station  
530 (Missoula, MT)
- 531 Perez B, Moreno JM (1998) Methods for quantifying fire severity in shrubland-fires. *Plant*  
532 *Ecology* **139**, 91-101.
- 533 Perez-Cabello F, Fernandez JR, Lloveria RM, Garcia-Martin A (2006) Mapping erosion-  
534 sensitive areas after wildfires using fieldwork, remote sensing, and geographic information  
535 systems techniques on a regional scale. *Journal of Geophysical Research--Biogeosciences*  
536 **111**, G04S10, doi:10.1029/2005JG000148
- 537 Peterson DL, Ryan KC (1986) Modeling postfire conifer mortality for long-range planning.  
538 *Environmental Management* **10**, 797-808.
- 539 Robichaud PR (2000) Fire effects on infiltration rates after prescribed fire in Northern Rocky  
540 Mountain forests, USA. *Journal of Hydrology* **231-232**, 220-229.
- 541 Robichaud PR, Beyers JL, Neary DG (2000) Evaluating the effectiveness of postfire  
542 rehabilitation treatments. USDA Forest Service, Rocky Mountain Research Station, General  
543 Technical Report RMRS-GTR-63 (Ogden, UT)
- 544 Robichaud PR, Elliot WJ, Pierson FB, Hall DE, Moffet CA, and Ashmum LF (2007a) Erosion  
545 risk management tool (ERMiT) user manual (version 2006.01.18). USDA Forest Service,  
546 Rocky Mountain Research Station, General Technical Report RMRS-GTR-188 (Ogden, UT)
- 547 Robichaud PR, Lewis SA, Laes DYM, Hudak AT, Kodaly RF, Zamudio JA (2007b) Postfire soil  
548 burn severity mapping with hyperspectral image unmixing. *Remote Sensing of Environment*.  
549 *In press*.

- 550 Rogan J, Franklin J (2001) Mapping wildfire burn severity in southern California forests and  
551 shrublands using enhanced thematic mapper imagery. *Geocarto International* **16**, 89-99.
- 552 Roldan-Zamarron A, Merino-de-Miguel S, Gonzalez-Alonso F, Garcia-Gigorro S, Cuevas JM  
553 (2006) Minas de Riotinto (south Spain) forest fire: Burned area assessment and fire severity  
554 mapping using Landsat 5-TM, Envisat-MERIS, and Terra-Modis postfire images. *Journal of*  
555 *Geophysical Research--Biogeosciences* **111**, G04S11, doi:10.1029/2005JG000136
- 556 Rothermel RC (1972) 'A mathematical model for predicting fire spread in wildland fuels.' USDA  
557 Forest Service, Intermountain Forest and Range Experiment Station, Research Paper INT-  
558 115 (Ogden, UT).
- 559 Roy DP, Boschetti L, Trigg SN (2006) Remote sensing of fire severity: assessing the  
560 performance of the normalized burn ratio. *IEEE Geoscience and Remote Sensing Letters* **3**,  
561 112-116.
- 562 Ruiz-Gallardo JR, Castano S, Calera A (2004) Application of remote sensing and GIS to locate  
563 priority intervention areas after wildland fires in Mediterranean systems: a case study from  
564 south-eastern Spain. *International Journal of Wildland Fire* **13**, 241-252.
- 565 Ryan KC (1981) 'Evaluation of a passive flame-height sensor to estimate forest fire intensity.'  
566 USDA Forest Service, Pacific Northwest Forest and Range Experiment Station, Research  
567 Note PN@-390 (Portland, OR).
- 568 Ryan KC (2002) Dynamic interactions between forest structure and fire behavior in boreal  
569 ecosystems. *Silva Fennica* **36**, 13-39.
- 570 Ryan KC, Frandsen WH (1991) Predicting postfire mortality of seven western conifers.  
571 *Canadian Journal of Forest Research* **18**, 1291-1297.
- 572 Ryan KC, Noste NV (1985) Evaluating prescribed fires. In 'Proceedings, symposium and

- 573 workshop on wilderness fire, 1983 November 15018, Missoula, MT'. (Eds JE Lotan, BM  
574 Kilgore, WC Fischer and RW Mutch) pp. 230-238. USDA Forest Service, Intermountain  
575 Forest and Range Experiment Station, General Technical Report INT-182 (Missoula, MT)
- 576 Sackett SS, Haase SM, and Harrington MG (1996) Lessons learned from fire use restoring  
577 southwestern ponderosa pine ecosystems. Pages 54-61 In 'Conference on adaptive ecosystem  
578 restoration and management: restoration of cordilleran conifer landscapes of North America'.  
579 (Eds WW Covington and PK Wagner) pp. 54-61. USDA Forest Service, General Technical  
580 Report RM-GTR-278, Rocky Mountain Forest and Range Experiment Station (Fort Collins,  
581 CO)
- 582 Salazar LA, Bradshaw LS (1986) Display and interpretation of fire behavior probabilities for  
583 long-term planning. *Environmental Management* **10**, 393-402.
- 584 Scott JH, Reinhardt ED (2001) 'Assessing crown fire potential by linking models of surface and  
585 crown fire behavior.' USDA Forest Service, Rocky Mountain Research Station, Research  
586 Paper RMRS-RP-29 (Fort Collins, CO)
- 587 Simard AJ (1991) Fire severity, changing scales, and how things hang together. *International*  
588 *Journal of Wildland Fire* **1**, 23-34.
- 589 Smith AMS, Wooster MJ, Drake NA, Dipotso FM, Falkowski MJ, Hudak AT (2005) Testing the  
590 potential of multi-spectral remote sensing for retrospectively estimating fire severity in  
591 African savannahs. *Remote Sensing of Environment* **97**, 92-115.
- 592 Stronach NH, McNaughton SJ (1989) Grassland fire dynamics in the Serengeti ecosystem, and a  
593 potential method of retrospectively estimating fire energy. *Journal of Applied Ecology* **26**,  
594 1025-1033.

- 595 Sugihara NG, van Wagtendonk, JW, Fites-Kaufman, J (2006) Fire as an ecological process. In  
596 'Fire in California's ecosystems'. (Eds NG Sugihara, JW van Wagtendonk, KE Shaffer, J  
597 Fites-Kaufman and AE Thode) pp. 58-74. (University of California: Los Angeles, CA)
- 598 Tolhurst KG (1995) Fire from a flora, fauna and soil perspective: sensible heat measurement.  
599 *CALM Science Supplement 4*, 45-88.
- 600 Turner MG, Hargrove WW, Gardner RH, Romme WH (1994) Effects of fire on landscape  
601 heterogeneity in Yellowstone National Park, Wyoming. *Journal of Vegetation Science 5*,  
602 731-742.
- 603 Turner MG, Romme WH, Gardner RH (1999) Prefire heterogeneity, fire severity, and early  
604 postfire plant reestablishment in subalpine forests of Yellowstone National Park, Wyoming.  
605 *International Journal of Wildland Fire 9*, 21-36.
- 606 Valette J-C, Gomendy V, Marechal J, Houssard C, Gillon D (1994) Heat transfer in the soil  
607 during very low-intensity experimental fires: the role of duff and soil moisture content.  
608 *International Journal of Wildland Fire 4*, 225-237.
- 609 van Wagner CE (1973) Height of crown scorch in forest fires. *Canadian Journal of Forest*  
610 *Research 3*, 373-378.
- 611 van Wagtendonk JW, Root RR, Key CH (2004) Comparison of AVIRIS and Landsat ETM+  
612 detection capabilities for burn severity. *Remote Sensing of Environment 92*, 397-408.
- 613 Vesk PA, Westoby M (2004) Sprouting ability across diverse disturbances and vegetation types  
614 worldwide. *Journal of Ecology 92*, 310-320.
- 615 Wade DD (1993) Thinning young loblolly pine stands with fire. *International Journal of*  
616 *Wildland Fire 3*, 169-178.

- 617 Wang GG, Kembell KJ (2003) The effect of fire severity on early development of understory  
618 vegetation following a stand replacing wildfire. In 'Second International wildland fire  
619 ecology and fire management congress and fifth symposium on fire and forest meteorology'  
620 pp. 3b.2:1-11. (American Meteorological Society: Boston, MA)
- 621 Weber RO (2001) Wildland fire spread models. In 'Forest fires. Behavior and ecological effects'.  
622 (Eds EA Johnson and K Miyanishi) pp. 151-169. (Academic Press: San Diego, CA)
- 623 Wells CG, DeBano LF, Lewis CE, Fredriksen RL, Franklin EC, Froelich RC, Dunn PH (1979)  
624 'Effects of fire on soil. A state-of-knowledge review.' USDA Forest Service, General  
625 Technical Report WO-7.
- 626 Whelan RJ (1995) 'The ecology of fire.' (Cambridge University Press: Cambridge, U.K.)
- 627 White JD, Ryan KC, Key CC, Running SW (1996) Remote sensing of forest fire severity and  
628 vegetation recovery. *International Journal of Wildland Fire* **6**, 125-136.
- 629 White PS, Pickett STA (1985) Natural disturbance and patch dynamics: an introduction. In 'The  
630 ecology of natural disturbance and patch dynamics'. (Eds STA Pickett and PS White) pp. 1-  
631 13. (Academic Press, Inc.: San Diego, CA)
- 632 Williams RJ, Gill AM, Moore PHR (1998) Seasonal changes in fire behaviour in a tropical  
633 savanna in northern Australia. *International Journal of Wildland Fire* **8**, 227-239.
- 634 Wilson CJ, Carey JW, Beeson PC, Gard MO, Lane LJ (2001) A GIS-based hillslope erosion and  
635 sediment delivery model and its application in the Cerro Grande burn area. *Hydrological  
636 Processes* **15**, 2995-3010.
- 637 Wooster MJ, Zhukov B, Oertel D (2003) Fire radiative energy for quantitative study of biomass  
638 burning: derivation from the BIRD experimental satellite and comparison to MODIS fire  
639 products. *Remote Sensing of Environment* **86**, 83-107.

640  
 641 **Table 1.** The matrix originally proposed by Ryan and Noste (1985) that related changes in  
 642 aboveground vegetation and soil organic matter has generally been simplified to a table such as  
 643 the below; modified from Ryan (2002) and Turner et al. (1994).  
 644

---

| Fire severity                       | Description   |
|-------------------------------------|---|
| Unburned                            | Plant parts green and unaltered, no direct effect from heat.  |
| Scorched                            | Unburned but plants exhibit leaf loss from radiated heat.   |
| Light                               | Canopy trees with green needles although stems scorched.<br>Surface litter, mosses, and herbs charred or consumed.<br>Soil organic layer largely intact and charring limited to a few mm depth.                             |
| Moderate or<br>severe surface burn: | Trees with some canopy cover killed, but needles not consumed.<br>All understory plants charred or consumed.<br>Fine dead twigs on soil surface consumed and logs charred.<br>Pre-fire soil organic layer largely consumed. |
| Deep burning or<br>crown fire:      | Canopy trees killed and needles consumed.<br>Surface litter of all sizes and soil organic layer largely consumed.<br>White ash deposition and charred organic matter to several cm depth.                                   |

---

645  
 646  
 647  
 648  
 649  
 650  
 651  
 652  
 653  
 654  
 655  
 656  
 657  
 658  
 659  
 660  
 661  
 662  
 663  
 664  
 665  
 666  
 667  
 668  
 669  
 670  
 671  
 672  
 673  
 674  
 675  
 676  
 677

678  
 679  
 680  
 681

**Table 2.** Summary of fire terminology and metrics

|                            | <b>Fire Intensity</b>   | <b>Fire Severity</b>  | <b>Burn Severity</b>  | <b>Ecosystem Responses</b>   |
|----------------------------|---|---|---|--|
| <b>Appropriate usage</b>   | Energy output from fire.  | Aboveground and below ground organic matter consumption from fire.  | Aboveground and below ground organic matter consumption from fire. Sometimes subdivided into ‘vegetation burn severity’ and ‘soil burn severity’  | Functional processes that are altered by fire including regeneration, recolonization by plants and animals and watershed hydrology parameters processes altered by fire.   |
| <b>Metrics</b>             | Strictly speaking is the time-averaged energy flux in Watt m <sup>-2</sup> , but more broadly can be measured as fireline intensity, temperature, residence time, radiant energy and other. | Aboveground measures include tree crown canopy scorch, crown volume kill, bole height scorch, skeleton twig diameter. Belowground and soil measures include ash deposition, surface organic matter, belowground organic matter contributing to soil structure, degree of hydrophobicity, and heat-induced oxidation of minerals.<br>Mortality is a common measure that is best applied to non-sprouting trees in surface fire regimes. In crown fire regimes aboveground mortality may be useful when fires are patchy. | Often used interchangeably with fire severity. Usually the term is applied to soils and designated ‘soil burn severity.’ In the U.S. it is the preferred term used in postfire BAER assessments and is considered to be the relative change due to fire; i.e., two soils with poor structure and low organic matter content may be rated differently if one was in that condition prior to the fire and another was not. Degree of severity may be influenced by socio-political concerns such as values at risk. | Vegetative regeneration, plant community composition and diversity, and plant and animal recolonization are important biotic parameters. Watershed hydrological processes such as dry ravel, erosion, and debris flows are the more important abiotic processes. |
| <b>Inappropriate usage</b> | Should never be used to describe fire effects such as those described under any of the remaining columns.   | Should not include ecosystem responses. Also, in shrubland ecosystems, complete above- and belowground mortality should not be considered here because it depends on vegetation composition and the proportion of sprouting and non-sprouting species.  | Should not include ecosystem responses. Also, this term should be restricted to field measurements and not be used to name remote sensing indices because the interpretation of remote data is dependent on ground-truthing with field measurements of burn severity; calling both measures burn severity is circular.  | Correlations between severity and ecosystem responses demonstrated in one system should not be considered universal for all ecosystems.  |

**Box 1.** Interpreting the Landsat dNBR signal in terms of fire severity and ecosystem response in crown-fire chaparral shrublands

In late October 2003 five large wildfires burned more than 200,000 ha in southern California. A total of 250 0.1-ha plots were sampled in these burned areas to assess fire severity and vegetation recovery (Keeley, Brennan and Pfaff, in preparation). Fire severity was assessed using the twig diameter method commonly used in crown fire ecosystems (Moreno and Oechel 1989; Perez and Moreno 1998) on multiple samples of the same shrub (*Adenostoma fasciculatum*) at all sites. Vegetation recovery was based on plant cover in the first spring following fires. The early assessment dNBR data were provided by EROS data center (USGS, Sioux Falls, SD).

The Landsat TM index is strongly correlated with our field measurement of fire severity (Fig. 3a), explaining over a third of the variation between these 250 sites. However, if dNBR is then used to predict ecosystem response variables we find little or no relationship. Total vegetative recovery (Fig. 3b) was very weakly related to dNBR and explained only about 1% of the variation, and there was no significant relationship with woody cover ( $P = 0.94$ , not shown), or percentage of the prefire *Adenostoma fasciculatum* population resprouting (Fig. 3c). These results argue against the concept of a composite burn index that mixes fire severity and ecosystem responses, even if such composites generate significant relationships with dNBR. For example, a standardized index that includes fire severity (Fig. 3a) and the two ecosystem impact variables (Figs. 3b, 3c) was created and it did generate a highly significant relationship with dNBR ( $P < 0.000$ ), but clearly this “composite index” is driven by the fire severity response variable (Fig. 3a).

Further complications arise with composite indices when adding in terms that have species-specific differences in the direction of response. For example, in this data set fire severity was slightly negatively correlated with log seedling recruitment of facultative-seeding shrubs, whereas fire severity was positively correlated with obligate seeding shrub recruitment. These shrublands may be an example in which remote sensing data can provide some information on fire severity but has limited predictive ability for ecosystem impacts, thus requiring coupling of remote sensing data with field studies (e.g., Ludwig et al. 2007).




Fig. 3

37

38 Figure Legends

39

40 Fig. 1. Schematic representation relating the energy output from a fire (fire intensity), the impact  
41 as measured by organic matter loss (fire severity), and ecosystem responses and societal impacts.

42

43 Fig. 2. (a) Arizona ponderosa pine forest illustrating different degrees of fire severity; entire  
44 scene burned, foreground mostly low severity with patches of scorched canopy of moderate  
45 severity and background high severity, b) soil burn severity assessment with characteristics of  
46 high severity, including heavy white ash deposition indicating loss of substantial levels of  
47 organic matter and loose unstructured soil, c) chaparral shrublands with large shrub skeletons  
48 retaining small twigs indicative of low fire severity and d) high fire severity.

49


50 Fig. 3. Relationship of Landsat TM differenced Normalized Burn Ratio based on spectral  
51 analysis of Landsat TM sensing data taken in the first growing season after the Fall 2003  
52 wildfires in southern California chaparral (scaled from 0 – 250) to (a) field measurement of fire  
53 severity and the extent to which dNBR can predict ecosystem response variables of (b) first year  
54 plant cover and (c) resprouting percentage of the common shrub *Adenostoma fasciculatum*, for  
55 250 sites distributed across the Otay, Cedar, Paradise, Old and Grand Prix fires (Landsat imagery  
56 from the USGS EROS Center, field data from Keeley, Brennan and Pfaff, in press).

57

58

59

60



61  
62  
  
63  
  
64  
  
65  
  
66  
  
67  
  
68  
  
69  
  
70  
  
71  
  
72  
  
73  
  
74  
  
75



76

77

78

79

80

81

82

83

84

85

86