

Climatic influences on fire regimes in montane forests of the southern Cascades, California, USA

A. H. Taylor^{A,D}, V. Trouet^{A,B} and C. N. Skinner^C

^ADepartment of Geography, The Pennsylvania State University, University Park, PA 16802, USA.

^BPresent address: Swiss Federal Institute, WSL, Zürcherstrasse 111, Birmensdorf CH8903, Switzerland.

^CUSDA Service Center, Pacific Southwest Research Station, 3644 Avtech Parkway, Redding, CA 96002, USA.

^DCorresponding author. Email: aht1@psu.edu

Abstract. The relationship between climate variability and fire extent was examined in montane and upper montane forests in the southern Cascades. Fire occurrence and extent were reconstructed for seven sites and related to measures of reconstructed climate for the period 1700 to 1900. The climate variables included the Palmer Drought Severity Index (PDSI), summer temperature (TEMP), NINO3, a measure of the El Niño–Southern Oscillation (ENSO), and the Pacific Decadal Oscillation (PDO). Fire extent at the site and regional scale was associated with dry and warm conditions in the year of the fire and regional fire extent was not associated with ENSO or PDO for the full period of analysis. The relationship between regional fire extent and climate was not stable over time. The associations of fire extent with PDSI and TEMP were only significant from ~1775 onward and the associations were strongest between 1805 and 1855. PDO and fire extent were also associated during the 1805–1855 period, and ENSO was associated with fire extent before 1800, but not after. The interannual and interdecadal variability of the fire response to temperature and drought suggests that increased periods of regional fire activity may occur when high interannual PDSI variation coincides with warm decades.

Additional keywords: American Pacific coast, climatic variation, El Niño–Southern Oscillation, fire ecology, Pacific Decadal Oscillation.

Introduction

Fire regimes in the forested landscapes of the southern Cascades are controlled by environmental processes and interactions that vary across a range of spatio-temporal scales (Skinner and Taylor 2006). For example, at local scales, spatial and temporal variation in fire frequency, extent, and severity before widespread Euro-American settlement is related to elevation, slope aspect, species composition, and time since the last fire (Taylor 2000; Beaty and Taylor 2001; Bekker and Taylor 2001; Taylor and Solem 2001; Taylor and Skinner 2003). However, at regional scales, spatial and temporal variation in fire regimes is linked to changes in land-use practices such as fire suppression and to climatic variability (Beaty and Taylor 2001; Taylor and Solem 2001; Norman and Taylor 2003; Skinner and Taylor 2006). Years of widespread burning in forests of Pacific Coast states before 20th century fire exclusion were broadly synchronous and usually associated with drought conditions (e.g. Swetnam and Baisan 2003; Hessl *et al.* 2004; Taylor and Beaty 2005), but drought may not be a precondition for widespread burning during some periods (Taylor and Beaty 2005). Thus, fire–climate interactions can be complex (e.g. Swetnam 1993; Grissino-Mayer and Swetnam 2000) and analyses of how climate modulates fire regimes over long periods are needed to understand fire–climate relationships and how fire regimes may respond to projected temperature increases associated with global climate change (Hayhoe *et al.* 2004). In the southern Cascades, where the present study was conducted, fire regimes are thought to be particularly sensitive to

temperature increases because of their effect on snowpack duration and fire season length (Dettinger and Cayan 1995; Lenihan *et al.* 2003; Hayhoe *et al.* 2004).

Interannual climatic variation along the Pacific Coast is strongly influenced by variation in sea surface temperatures (SST) in the eastern and central tropical Pacific Ocean associated with the El Niño–Southern Oscillation (ENSO). ENSO is a coupled ocean–atmospheric process that generates regionally distinct high frequency (3–7 year) variations in temperature and precipitation (Diaz and Markgraf 2000). Conditions in the American South-west (SW) during a warm phase ENSO (El Niño) are typically wet and warm, whereas during a cool phase ENSO (La Niña), conditions are cooler and drier. The pattern of climatic variation associated with ENSO is the opposite in the American Pacific North-west (PNW). SST variation associated with the Pacific Decadal Oscillation (PDO) in the north Pacific generates ‘ENSO-like’ climatic variation at interdecadal time scales (Dettinger *et al.* 2000). When the PDO is in a warm phase, conditions in the SW are wetter than average, and they are dry and warmer in the PNW; the opposite pattern prevails during a cool-phase PDO (Nigam *et al.* 1999). Moreover, the PDO interacts with ENSO, modulating the amplitude and geographic expression of climatic variation associated with ENSO (Gershunov *et al.* 1999). The southern Cascades are located in the pivot zone of the ENSO–PDO precipitation dipole that is located at ~40–45°N and that shifts north or south on interannual and decadal time scales (Dettinger *et al.* 1998). Thus, interannual

and interdecadal fire–climate associations in this location may be similar to those in the PNW in some periods and to the SW in others (Westerling and Swetnam 2003; Taylor and Beaty 2005).

The goal of the present study is to identify the relationships between climate variability (ENSO, PDO, drought) and fire occurrence and extent during the pre-fire suppression period in the southern Cascades at interannual to multidecadal time scales. In our analysis, we place particular emphasis on identifying fire–climate relationships: (1) across the range of dominant forest types in the montane and upper montane zones in the southern Cascades; and (2) during different time periods prior to the establishment of organised fire suppression in 1905 (Strong 1973). Specifically, we address the following questions: (1) How have fire extent and frequency varied over time? (2) Were widespread fires synchronous; did they burn primarily in dry years and were years with less burning wet ones? (3) Has fire frequency and extent varied with ENSO, PDO and their interactions? and (4) Have fire–climate relationships varied over time or have they remained constant?

Materials and methods

Study area

Montane and upper montane forests in the southern Cascades are segregated primarily by elevation and secondarily by topographic position (Barbour 1988; Parker 1991). Low elevation (<600 m) xeric sites on the west side of the range are usually dominated by ponderosa pine (*Pinus ponderosa*) with black oak (*Quercus kelloggii*) as an important associate. On the east side of the range, low-elevation (<1700 m) xeric uplands are dominated either by Jeffrey pine (*Pinus jeffreyi*) or ponderosa pine, and western juniper (*Juniperus occidentalis* ssp. *occidentalis*) may be an associate on the driest sites. Mixed conifer forests dominate the landscape above the lower montane zone to ~2000 m. In this zone, any of six conifer species – ponderosa pine, Douglas-fir (*Pseudotsuga menziesii*), sugar pine (*Pinus lambertiana*), incense cedar (*Calocedrus decurrens*), Jeffrey pine, and white fir (*Abies concolor*) – may co-occur and share dominance in a stand depending on site conditions and stand history. A subcanopy of the hardwoods black oak, Pacific madrone (*Arbutus menziesii*), big leaf maple (*Acer macrophyllum*), Pacific dogwood (*Cornus nuttallii*), and canyon live oak (*Quercus chrysolepis*) may be present in stands found on the more temperate west side of the range. In the mixed conifer zone, the tree cover is often interrupted by stands of montane chaparral. Chaparral shrubs are usually <2 m tall and the most common species are green-leaf manzanita (*Arctostaphylos patula*), California lilac (*Ceanothus* sp.), and shrub oaks (*Quercus* sp.). Montane chaparral shrubs are fire-adapted (e.g. sprout following fires or regenerate from fire-stimulated seeds) and chaparral appears to occupy sites that once experienced high-severity fire or are too poor to support trees (Wilken 1967; Nagel and Taylor 2005). Mixed white fir and red fir (*Abies magnifica* var. *shastensis*) forests occur above the mixed conifer zone. Above 2000 m, white fir is replaced by red fir and western white pine (*Pinus monticola*), which are upper montane zonal dominants. Topographic lowlands that receive cold air drainage are dominated by either Jeffrey pine or lodgepole pine (*Pinus contorta*). In these locations, lodgepole pine is more common on the more mesic sites or sites that have experienced

high-severity fire. Forests and woodlands above 2400 m are dominated by mountain hemlock (*Tsuga mertensiana*) or whitebark pine (*Pinus albicaulis*) (Barbour 1988; Parker 1991).

Fire occurrence and extent were studied at seven sites that include the dominant montane and upper montane forest types found in the southern Cascades (Fig. 1; Table 1). Elevations for sites ranged from 1136 to 2646 m and study areas ranged from 2 to 26 km², except for Lassen Meadows (LM), which covered 700 km² (Table 1). Climate in the southern Cascades is characterised by warm, dry summers and cold, wet winters. Annual precipitation decreases from west (Mineral, 137 cm; elevation 1486 m) to east (Susanville, 36 cm; elevation 1300 m) and is generally lower north of Mount Shasta on either side of the range owing to the rain shadow of the Klamath Mountains. Most (>75%) precipitation falls as snow between November and April. Depth of April snowpack in the upper montane zone commonly exceeds 2 m on the west side of the range where snow may persist into mid–late June in wet years (Taylor 1990). Mean monthly temperatures in Mineral on the west side of the range are lowest in January (–1.1°C) and highest in July (16.9°C). The snowpack on the east side of the range is usually shallower and temperatures are more continental. Mean monthly temperatures for Mount Hebron at 1295 m in January are –2.7°C and 17.1°C in July (Skinner and Taylor 2006). Additional details on site conditions are provided in references cited in Table 1.

Human influence on fire regimes in the southern Cascades has varied over time. Native Americans are known to have used fire to collect insects, to drive game and to encourage certain plants used for food and fibre (Schulz 1954). Euro-American settlement began ~1850. Sheep and cattle grazing, which influences the continuity of grassy fuels, began in the 1860s (Norman and Taylor 2005). A policy of suppressing fire was first implemented along railroad lines in the 1880s (Skinner and Taylor 2006) and became more widespread with establishment of the Forest Reserve system in 1905 (Strong 1973; Norman and Taylor 2005).

Fire record

Fire occurrence and extent during the pre-fire suppression period were reconstructed using fire-scarred cross-sections removed from fire-scarred stumps, logs, and trees (Arno and Sneek 1977). Samples were widely distributed in each study area to capture spatial variability in fire regimes related to environmental and species compositional gradients (Table 1). The number of fire scar samples per site ranged from 39 to 152; a total of 530 samples are included in the current study (Table 1). Cross-sections were sanded to a high polish and each sampled tree-ring series was crossdated using standard dendrochronological techniques (Stokes and Smiley 1968). The calendar year of a tree-ring with a fire scar lesion in it was then recorded as the fire date. Fire scars that could not be crossdated were not used in subsequent analyses.

Fire regimes were characterised for each site and for the southern Cascades region (all sites). For each site, we identified fire years recorded by any, 10% or more, and 25% or more of samples. For the southern Cascades region, we first identified fire years that were recorded by two or more samples at each site and used these fire dates for characterising regional




Fig. 1. Location of study sites with fire scar samples in the southern Cascade Range, California. National Forest and National Park lands are shown in light and dark grey, respectively.

Table 1. Location and sample characteristics for sites in the southern Cascades

The forest types are: ponderosa pine (PP), ponderosa pine–Jeffrey pine (PP-JP), white fir–Jeffrey pine (WF-JP), mixed conifer (MC), red fir–white fir (RF-WF), red fir–western white pine (RF-WWP), lodgepole pine (LP), red fir–mountain hemlock (RF-MH). Additional details on forest conditions and fire regimes in each area are given in the cited reference

Site	Elevation (m)	Area (km ²)	Samples	Time period	Reference	Forest dominant
Lava Beds (LB)	1580–1670	13.7	63	1563–1904	Miller and Heyerdahl (in press)	PP
Goosenest (GAMA)	1495–1800	2	44	1375–1935	Ritchie (2005)	MC
Thousand Lakes (TLW)	1700–2646	20.4	50	1652–1942	Bekker and Taylor (2001)	RF-MH/LP/RF-WF/WF-JP
Lassen Meadows (LM)	1650–2300	700	152	1520–1944	Norman and Taylor (2003)	PP
Prospect Peak (PP)	1800–2420	26.3	126	1507–1937	Taylor (2000)	RF-WWP/WF-JP/JP-PP
Caribou (CAR)	2060–2390	9.5	39	1735–1982	Taylor and Solem (2001)	LP/RF-WWP/WF-JP
Cub Creek (CCRNA)	1136–2044	15.9	56	1616–1926	Beatty and Taylor (2001)	MC

fire extent. An index of variability in regional annual fire extent was calculated by first determining the percentage of samples that recorded a fire in each year at each site. We then summed the site percentages as our index of annual fire extent for the

southern Cascade region (Fig. 2). This index provided a measure of how widespread burning was both within and among sites, but no assumptions were made about the number or spread of fires represented by the index.




Fig. 2. Proportion of samples scarred in each study area and cumulative proportion for all sites in the southern Cascades, California. *n*, number of fire scar samples.

Climate record

Four tree-ring-based reconstructions of climatic variables were used to evaluate the relationships between climate and fire. First, we used the Palmer Drought Severity Index (PDSI) as an index of drought (Palmer 1965). PDSI is a composite index that integrates immediate and lagged precipitation and temperature in estimating drought severity. We used the reconstructed summer

PDSI at grid point 35 to represent drought conditions in the southern Cascades (Cook and Krusic 2004). Second, as an index of temperature we used the reconstruction of western North American summer temperature (TEMP) by Briffa *et al.* (1992) (gridpoint 16), which expresses temperature variation relative to the 1951–70 base period. Third, we used a reconstruction of NINO3, a measure of tropical Pacific SST temperature variation

that is frequently used as an ENSO index (Cook 2000). Finally, a reconstruction developed by Gedalof *et al.* (2002) was used as our index of the PDO.

Fire–climate analysis

We used a combination of correlation analysis and superposed epoch analysis (SEA) to examine interannual and interdecadal relationships between TEMP, PDSI, ENSO, PDO, and fire (Haurwitz and Brier 1981; Baisan and Swetnam 1990). For the interannual analysis, SEA was used to evaluate the relationship between events (fire years) and climate, by superposing a window of contemporaneous and lagged climatic conditions over each event year. Significance levels between events and climatic conditions were determined from bootstrapped confidence interval estimates (95%) based on Monte Carlo simulations (Mooney and Duval 1993). SEA was performed separately for each site for fire years of different fire extent (any, $\geq 10\%$ of samples) and for years of different fire extent in the region (≥ 1 , ≥ 2 or ≥ 3 sites), and for non-fire years, to identify climatic conditions conducive to and unfavourable for fire.

Correlation analysis was also used to identify the association between interannual variability in fire extent and climate. For each site, and for the region, we calculated the Pearson product moment correlation coefficient of fire extent with each climate variable for the period 1700–1900. We used first differences (value (year t) – value (year $t - 1$)) for each variable to calculate correlation coefficients to emphasise the interannual variability in each time series.

Decadal-scale relationships between fire and climate were identified using correlation analysis. Pearson product moment correlation coefficients were calculated for fire and climate time series of sequential 21-year non-overlapping means (Swetnam 1993). We extended the time series for the decadal analysis back to 1600 to increase sample size ($n = 10$ to $n = 15$) for calculation of the correlation coefficients. Four of the seven sites had a record of fire back to 1600.

Results

Fire record and synchronicity of fire

The southern Cascade forests have a long history of fire. The record of fire spans the period 1375 to 1942 and the number of fire dates per site ranges from 23 to 197 (Table 1). We chose the period 1700–1900 for the climate analysis during which all sites and a minimum of 100 samples recorded fire (Fig. 2).

Fire dates co-occurred among sites much more frequently than expected by chance alone ($\chi^2 = 69.2, P < 0.001$). For example, over a 200-year period, seven fire dates are expected to co-occur among three or more sites; our record had 29 such co-occurring fire dates. Strong temporal synchrony in fire dates among widely dispersed sites suggests a strong regional influence on the occurrence and extent of fire in the southern Cascades.

Fire–climate interactions

Interannual relationships

Results of the SEA and correlation analysis of fire and PDSI for individual sites were similar for the 1700–1900 period (Fig. 3;

Table 2). For SEA, fire years were associated with low moisture conditions and the opposite was true for non-fire years, except in CAR and LB where fire years and PDSI were unrelated (Fig. 3; Table 2). Larger burns ($\geq 10\%$ and $\geq 25\%$) occurred during drier years, and antecedent climatic conditions (wet) were associated with fire years only in GAMA. For the correlation analysis, variation in fire extent was associated with dry and warm summer conditions in the year of the fire (Table 2). Variation in fire extent was also positively correlated with NINO3 in three of the seven sites; no correlation was found between fire extent and PDO at any of the sites.

Fire extent among sites was also strongly associated with drought. The SEA of PDSI indicated fire years are associated with low moisture conditions, with more sites burning during the driest years, and non-fire years are wet (Fig. 4). The SEA showed no association between regional fire extent and ENSO, PDO, or TEMP. However, correlation analysis indicated a strong association between dry conditions, high summer temperature and regional fire extent (Fig. 5a; Table 2). Regional fire extent was not correlated with ENSO or PDO.

Temporal variability

The relationship between regional fire extent, PDSI, and TEMP was not consistent over time. Correlation coefficients for first differences of fire extent with PDSI and TEMP were only significant from ~ 1775 onward and the correlations were strongest between 1805 and 1855 (Fig. 5b). Although there was no correlation between fire extent and PDO for the 1700–1900 period (Table 2), PDO and fire extent were positively correlated for the 1805–1855 period. Fire extent and ENSO were positively correlated before 1800, but not after (Fig. 5b).

Temporal variation in the relationship between PDSI and regional fire extent was confirmed by separate SEA for the 1700–1800 and 1800–1900 time periods (Fig. 4). Widespread fires were associated with drought after but not before 1800. Moreover, non-fire years after 1800 were wet, but this was not the case before 1800. SEA showed no differences or significant relationships between fire extent, TEMP, ENSO, or PDO.

Interdecadal relationships

There was no significant correlation between regional fire extent and PDSI on a decadal time scale. However, variation in fire extent was correlated with variation in TEMP ($r = 0.45, P < 0.01$) and ENSO ($r = 0.48, P < 0.01$) (Fig. 6).

Discussion and conclusions

Fire occurrence in the southern Cascades was strongly synchronous despite long distances between sites and large differences in forest composition and elevation, which are variables known to be locally associated with variation in fire frequency, fire extent, and fire severity in the southern Cascades (Taylor 2000; Bekker and Taylor 2001; Skinner and Taylor 2006). The expected frequency of co-occurring fire for six of seven sites is once every 13 000 years. Yet six sites burned in the same year five times (1751, 1781, 1800, 1829, 1883) during our 200-year study period. This high synchrony of fire dates among sites is probably not caused by burns that spread continuously between sites. Instead, the synchrony reflects a strong

influence of regional climate that promotes conditions that are conducive to burning across the range of montane and upper montane forests present in the southern Cascades (Swetnam and Betancourt 1998; Donnegan *et al.* 2001).

There is a strong relationship between drought and widespread burning during the pre-fire suppression period in coniferous forests in the SW (Swetnam and Baisan 2003), PNW

(Heyerdahl *et al.* 2001; Hessl *et al.* 2004) and Rocky Mountain (RM; Donnegan *et al.* 2001) regions in the western USA. The overall effect of drought on temporal patterns of burning in the southern Cascades is similar. Widespread burning within the region was associated with both low moisture conditions and warm summer temperatures during the fire year. The opposite climatic conditions characterised non-fire years. Moreover,




Fig. 3. Superposed Epoch Analysis (SEA) of reconstructed Palmer Drought Severity Index (PDSI) (Cook and Krusic 2004) with non-fire years and fire years of different extent (i.e. at least one sample, $\geq 10\%$, and $\geq 25\%$ of samples scarred) in the southern Cascades, California, for the period 1700–1900. The analysis window includes up to 5 years before and 2 years after each fire or non-fire year. Values with filled symbols were statistically significant ($P < 0.05$).

across all sites, years of widespread burning did not lag years of high moisture availability. This suggests that fires are dependent on drought and warm summer temperatures and not increased production of fine fuels, at least across sites and forest types in the region. However, in the dry mixed conifer forests at GAMA, and in dry pine forests in the northern Sierra Nevada (Taylor and Beaty 2005), wet conditions preceded years with widespread fire. Presumably, increased production of fine grass and herbaceous fuels in wet years increased fuel connectivity, promoting widespread burning in subsequent dry years. This fire–climate pattern has also been documented for dry pine forests in the SW (Swetnam and Baisan 2003), northern Mexico (Stephens *et al.* 2003), and the RM (Veblen *et al.* 2000), but not in the PNW. In the dry pine forests of the PNW, increased growth of fine fuel in wet years is not necessary to precondition the landscape for widespread burning (Heyerdahl *et al.* 2001; Hessl *et al.* 2004).

Interannual variation in ENSO has been identified as a strong control on variation in fire extent in pine-dominated forests in the SW (Swetnam and Betancourt 1990), northern Mexico (Stephens *et al.* 2003), the PNW (Heyerdahl *et al.* 2002), and the RM (Veblen *et al.* 2000). In each of these regions, years with widespread burning are associated with the ENSO phase that corresponds with drier and warmer conditions. In the southern Cascades, ENSO was not a strong driver of variation in fire extent at the regional scale, though there was a weak influence

Table 2. Pearson correlation coefficients of fire extent and climate for the period 1700–1900 for seven sites and a composite of all sites, southern Cascade Range, California

The climatic variables are Palmer Drought Severity Index (PDSI), summer temperatures (TEMP), the El Niño–Southern Oscillation (ENSO), and the Pacific Decadal Oscillation (PDO); *, $P < 0.05$ or **, $P < 0.01$. See Table 1 for site definitions

Site	PDSI	TEMP	ENSO	PDO
LB	−0.06	−0.023	0.057	−0.01
GAMA	−0.208**	0.163*	−0.05	−0.01
TLW	−0.251**	0.171*	0.157*	0.07
LM	−0.206**	0.145*	0.153*	0.08
PP	−0.228**	0.192**	0.155*	0.03
CAR	−0.185**	0.07	−0.049	0.03
CCRNA	−0.2**	0.222**	0.04	0.10
Composite	−0.356**	0.252**	0.129	0.08

of ENSO on the extent of fires in three of the seven sites. Spatial variation in the position of zonal precipitation in the ENSO pivot zone on interannual time scales may reduce the coherency of the ENSO signal on fire climate in this region. In the northern Sierra Nevada, interannual variation in instrumental precipitation tends to be independent of ENSO (Schonher and Nicholson 1989).

Fire occurrence and extent in the western USA has been shown to vary with the PDO. In the southern RM, widespread fires in high-elevation forests are associated with a negative (cool phase) PDO (Schoennagel *et al.* 2005; Sibold and Veblen 2006). However, in lower-elevation pine forests in the PNW, widespread burning tended to coincide with a positive PDO (Hessl *et al.* 2004), where the PDO is a driver of multidecadal winter precipitation (Mantua *et al.* 1997). In some dry pine and mixed conifer forests in the northern Sierra Nevada, years with widespread burning are also associated with a negative PDO whereas non-fire years are associated with the opposite or positive PDO (Taylor and Beaty 2005; Moody *et al.* 2006). In the southern Cascades, variation in the frequency and extent of burning was not related simply to variation in the PDO. Instead, the relationship between fire extent and PDO varied by time period.

The sensitivity of fire activity in the southern Cascades to climatic variation was not consistent over time. There was a sharp peak in the strength of the association between concurrent low PDSI, warm summer temperatures and widespread burning in the ~1800–1850 period. One explanation for the peak during this 50-year period could be methodological, related to the large amplitude switch in PDSI and fire extent between the years 1829 and 1830. Periods with high amplitude changes can inflate running correlation coefficients (Swetnam and Betancourt 1998). However, the frequency-based SEA analysis confirms that drought was associated with fire during the 1800–1900 period, but not during the 1700–1800 period. Similarly, concurrent drought and fire extent in dry pine forests in the northern Sierra Nevada are associated during the 1775–1850 period, but not between 1700 and 1775 (Taylor and Beaty 2005). This suggests that the temporal change in fire–climate relationships affected a wide area.

The late 18th–early 19th century transition period (LEENT) has previously been identified as a period of shifting fire regimes and shifting fire–climate relationships in forests and woodlands in other parts of the western USA, north-western Mexico (Stephens *et al.* 2003), and even Argentina (Grissino-Mayer and




Fig. 4. Superposed epoch analysis (SEA) of reconstructed Palmer Drought Severity Index (PDSI) with non-fire years and fire years of different extent (≥ 1 , ≥ 2 or ≥ 3 sites) in the southern Cascade Range, California, for the periods 1700–1800 and 1800–1900. There were too few fires to conduct an SEA for burns of larger extent (four or more sites burned). The analysis window includes up to 5 years before and 2 years after each fire or non-fire year. Values with filled symbols were statistically significant ($P < 0.05$).




Fig. 5. (a) Pearson product moment correlation coefficients of first differences for regional fire occurrence and Palmer Drought Severity Index (PDSI) and (b) 50-year running correlation, plotted on the 26th year of the period, between fire occurrence, PDSI, summer temperature (TEMP), El Niño–Southern Oscillation (ENSO), and Pacific Decadal Oscillation (PDO). The PDSI values and corresponding correlation coefficients were inverted for presentation.

Swetnam 2000; Kitzberger *et al.* 2001). At many sites in the SW and north-western Mexico, this period was characterised by reduced fire occurrence, and a related increase in the synchrony of fire (Swetnam and Betancourt 1998; Grissino-Mayer and Swetnam 2000; Stephens *et al.* 2003; Swetnam and Baisan 2003; Sakulich and Taylor 2007). Some of these fire regime shifts have also been documented for forests in the RM (Veblen *et al.* 2000; Donnegan *et al.* 2001) and the PNW (Heyerdahl *et al.* 2002) during the same period. The hemispheric scale of the LEENT fire regime shift suggests a global-scale driver. A decadal-scale decline in ENSO frequency and amplitude during

the LEENT (Anderson *et al.* 1992; Cleaveland *et al.* 1992; Mann *et al.* 2000), in particular, and the corresponding dampening of dry and wet patterns, has been suggested as a driver of the fire regime shifts (Kitzberger *et al.* 2001; Stephens *et al.* 2003; Swetnam and Baisan 2003). The weak influence of ENSO on fire regimes in northern California may explain why no shifts in the frequency or synchrony of fire were observed during the LEENT in our study area.

The dampening of the ENSO signal is possibly linked to a shift of PDO during the LEENT to an unusually cool phase (D’Arrigo *et al.* 2001), as the PDO phase affects the strength of




Fig. 6. Interdecadal variation in regional fire extent, Palmer Drought Severity Index (PDSI), summer temperature (TEMP), and El Niño–Southern Oscillation (ENSO) (NINO3). All series were computed from 20-year, non-overlapping means (sums in the case of fire extent) and were smoothed using a cubic spline for presentation.

ENSO climatic effects (Gershunov and Barnett 1998; McCabe and Dettinger 1999). PDO and fire activity in the southern Cascades in the LEENT period are positively associated, which corresponds with a strengthening of the PDO teleconnection in northern California. The strengthening of the interannual association between drought, summer temperature, and fire during the LEENT in the southern Cascades and northern Sierra Nevada (Taylor and Beaty 2005), however, contrasts sharply with the decline in the responses of fire to interannual climate patterns found for the same period in the SW (Grissino-Mayer and Swetnam 2000; Swetnam and Baisan 2003). The LEENT generally corresponds to an extraordinarily cold period over the northern Hemisphere (Mann *et al.* 1998), but moisture conditions vary by region. Whereas conditions tended to be wetter than normal in California (Earle 1993) and the PNW (Graumlich 1987), the SW was characterised by an increasing frequency of dry springs and summers (Swetnam and Baisan 2003).

The cool and wet conditions in northern California during LEENT probably enhanced the interannual relationship between drought and fire extent. Severe drought is a prerequisite for fire occurrence and spread during conditions that are wet and cool compared with drier periods, especially across a range of montane and upper montane forest types where fuel is not an important limitation on fire spread (Swetnam and Betancourt 1990). Snowpacks persist longer in wet and cool years in the montane and upper montane forests of the southern Cascades, shortening fire season length (Taylor 1995, 2000). During a multidecadal period of cool–wet conditions, high summer temperatures would promote earlier snowmelt, earlier fuel moisture depletion, and a longer fire season – all conditions more favourable to fire spread.

The strengthening of fire–climate relationships during LEENT could also result from a change in the influence of Native

American burning on fire regimes. Native American tribes in the southern Cascades are known to have used fire to improve wildlife habitat, drive game, encourage certain plants for food and fibre, and to collect grasshoppers (Schulz 1954). Native American population in the region declined rapidly in the 19th century after contact with Euro-Americans (Cook 1943). However, the decline in Native American populations in the southern Cascades is thought to have occurred several decades after the onset of LEENT and the interhemispheric nature of the timing of altered fire regimes suggests a global and not local driver of fire regime change.

Fire occurrence and extent in the southern Cascades were also affected by interdecadal variation in ENSO and summer temperatures. Decades in which the El Niño phase dominated coincided with widespread fires and vice versa. Similarly, decades with higher summer temperatures experienced more widespread burning than cooler decades. The decadal-scale sensitivity of fire regimes to the ENSO cycle and summer temperature may be a long-term influence of ENSO and temperature on fuel production. Decadal time-scale sensitivity of fire regimes has been attributed to long-term temperature-related shifts in fuel production (Swetnam 1993) and to concentration of temperature variability in decadal frequencies (Swetnam and Betancourt 1998). Variation in fire extent, however, was not related to PDSI on an interdecadal scale, which stresses the importance for fire spread of high frequency (interannual) wet–dry cycles and the abrupt fuel moisture changes related to them (Swetnam 1993). The frequency-dependent fire response to temperature and drought suggests that regional fire activity is most extensive when high interannual variation in PDSI coincides with warm decades (Swetnam and Betancourt 1998).

Climate strongly influenced fire regimes in the southern Cascades during the pre-fire suppression period. Both interannual

and interdecadal climatic variation promoted conditions conducive to fire, but fire–climate relationships were not stable over time. The timing of changes in fire–climate relationships is similar to the timing of changes identified in other regions in both the northern and southern hemisphere (Kitzberger *et al.* 2001), suggesting a strong linkage between global circulation and climatic conditions conducive to fire.

The strong relationships between climate and pre-fire suppression fire regimes in the southern Cascades, particularly the association of fire with interdecadal temperature variations suggests that future climatic change is likely to affect fire regimes in the region. Climate models project warming for the southern Cascades with increased concentrations of greenhouse gases in the atmosphere (Hayhoe *et al.* 2004). Although active fire suppression and forest management have led to less overall area being burned than was likely in the past (Skinner and Chang 1996), 20th-century fire extent in the western United States remains strongly linked to climatic conditions (Westerling and Swetnam 2003; Trouet *et al.* 2006; Westerling *et al.* 2006). In fact, it may be that the relationship between climate and fire extent has been enhanced owing to efficiency of fire suppression forces controlling most fires while quite small except for those under more extreme conditions. However, because of anthropogenic interference on the one hand and the complex character of fire–climate interactions on the other, predicting which years will have big fires remains a challenge.

Acknowledgements

The present research was supported by the Lassen National Forest, cooperative agreements (02-CA-11272162–052, 04-JV-11272162–407) between the USDA Forest Service and the Pennsylvania State University, the interagency Joint Fire Sciences Program, and a George S. Deike Research Grant.

References

- Anderson RY, Lindsay J, Parker D (1992) Long-term changes in the frequency of occurrence of El Niño events. In 'El Niño: Historical and Paleoclimatic Aspects of the Southern Oscillation'. (Eds HF Diaz, V Markgraf) pp. 193–200. (Cambridge University Press: Cambridge)
- Arno SF, Sneek KM (1977) A method for determining fire history in coniferous forest of the mountain west. USDA Forest Service, General Technical Report INT-GTR-42. (Ogden, UT)
- Baisan CH, Swetnam TW (1990) Fire history on a desert mountain range: Rincon Mountain Wilderness, Arizona, USA. *Canadian Journal of Forest Research* **20**, 1559–1569. doi:10.1139/X90-208
- Barbour MG (1988) California upland forests and woodlands. In 'North American Terrestrial Vegetation'. (Eds MG Barbour, WD Billings) pp. 131–164. (Cambridge University Press: New York)
- Beaty RM, Taylor AH (2001) Spatial and temporal variation of fire regimes in a mixed conifer forest landscape, Southern Cascades, California, USA. *Journal of Biogeography* **28**, 955–966. doi:10.1046/J.1365-2699.2001.00591.X
- Bekker MF, Taylor AH (2001) Gradient analysis of fire regimes in montane forests of the southern Cascade range, Thousand Lakes Wilderness, California, USA. *Plant Ecology* **155**, 15–28. doi:10.1023/A:1013263212092
- Briffa KR, Jones PD, Schweingruber FH (1992) Tree-ring density reconstructions of summer temperature patterns across western North America since 1600. *Journal of Climate* **5**, 735–754. doi:10.1175/1520-0442(1992)005<0735:TRDROS>2.0.CO;2
- Cleaveland MK, Cook ER, Stahle DW (1992) Secular variability of the Southern Oscillation detected in tree ring data from Mexico and the southern United States. In 'El Niño: historical and paleoclimatic aspects of the Southern Oscillation'. (Eds HF Diaz, V Markgraf) pp. 271–291. (Cambridge University Press: Cambridge)
- Cook ER (2000) 'Niño 3 Index Reconstruction.' International Tree-Ring Data Bank. IGBP PAGES/World Data Center-A for Paleoclimatology. Data Contribution Series 2000–052. (NOAA/NGDC Paleoclimatology Program: Boulder, CO)
- Cook ER, Krusic PJ (2004) 'The North American drought atlas.' (Lamont-Doherty Earth Observatory and the National Science Foundation: New York)
- Cook SF (1943) The conflict between the California Indian and white civilization. In 'Green versus gold: sources in California's environmental history'. (Ed. C Merchant) pp. 55–59. (Island Press: Washington, DC)
- D'Arrigo RD, Villalba R, Wiles G (2001) Tree-ring estimates of Pacific decadal climate variability. *Climate Dynamics* **18**, 219–224. doi:10.1007/S003820100177
- Dettinger MD, Cayan DR (1995) Large-scale atmospheric forcing of recent trends toward early snowmelt runoff in California. *Journal of Climate* **8**, 606–623. doi:10.1175/1520-0442(1995)008<0606:LSAFOR>2.0.CO;2
- Dettinger MD, Cayan DR, Diaz HF, Meko MD (1998) North–south precipitation patterns in western North America on interannual-to-decadal time scales. *Journal of Climate* **11**, 3095–3111. doi:10.1175/1520-0442(1998)011<3095:NSPPIW>2.0.CO;2
- Dettinger MD, Cayan DR, McCabe GJ, Marengo JA (2000) Multiscale hydrologic variability associated with El Niño/Southern Oscillation. In 'El Niño and the Southern Oscillation: multiscale variability and global and regional impacts'. (Eds HF Diaz, V Markgraf) pp. 113–146. (Cambridge University Press: Cambridge)
- Diaz HF, Markgraf V (Eds) (2000) 'El Niño and the Southern Oscillation: multiscale variability and global and regional impacts.' (Cambridge University Press: Cambridge)
- Donnegan JA, Veblen TT, Sibold JS (2001) Climatic and human influences on fire history in Pike National Forest, central Colorado. *Canadian Journal of Forest Research* **31**, 1526–1539. doi:10.1139/CJFR-31-9-1526
- Earle CJ (1993) Asynchronous droughts in California streamflow as reconstructed from tree rings. *Quaternary Research* **39**, 290–299. doi:10.1006/QRES.1993.1036
- Gedalof Z, Mantua NJ, Peterson DL (2002) A multi-century perspective of variability in the Pacific Decadal Oscillation: new insights from tree rings and coral. *Geophysical Research Letters* **29**, 2204–2207. doi:10.1029/2002GL015824
- Gershunov A, Barnett T (1998) Interdecadal modulation of ENSO teleconnections. *Bulletin of the American Meteorological Society* **79**, 2715–2725. doi:10.1175/1520-0477(1998)079<2715:IMOET>2.0.CO;2
- Gershunov A, Barnett TP, Cayan DR (1999) North Pacific Interdecadal Oscillation seen as a factor in ENSO-related North American climate anomalies. *EOS, Transactions, American Geophysical Union* **80**, 25–36. doi:10.1029/99EO00019
- Graumlich LJ (1987) Precipitation variation in the Pacific North-west (1675–1975) as reconstructed from tree rings. *Annals of the Association of American Geographers. Association of American Geographers* **77**, 19–29. doi:10.1111/J.1467-8306.1987.TB00142.X
- Grissino-Mayer HD, Swetnam TW (2000) Century-scale climate forcing of fire regimes in the American South-west. *The Holocene* **10**, 213–220. doi:10.1191/095968300668451235
- Haurwitz MW, Brier GW (1981) A critique of the superposed epoch analysis method: its application to solar–weather relations. *Monthly Weather Review* **109**, 2074–2079. doi:10.1175/1520-0493(1981)109<2074:ACOTSE>2.0.CO;2
- Hayhoe K, Cayan D, Field CB, Frumhoff PC, Maurer EP, *et al.* (2004) Emissions pathways, climate change, and impacts on California. *Proceedings*

- of the National Academy of Sciences of the United States of America* **101**, 12422–12427. doi:10.1073/PNAS.0404500101
- Hessl AE, McKenzie D, Schellhaas R (2004) Drought and Pacific Decadal Oscillation linked to fire occurrence in the inland Pacific North-west. *Ecological Applications* **14**, 425–442. doi:10.1890/03-5019
- Heyerdahl EK, Brubaker LB, Agee JK (2001) Spatial controls of historical fire regimes: a multi-scale example for the Interior West, USA. *Ecology* **82**, 660–678.
- Heyerdahl EK, Brubaker LB, Agee JK (2002) Annual and decadal climate forcing of historical fire regimes in the interior Pacific Northwest. *The Holocene* **12**, 597–604.
- Kitzberger T, Swetnam TW, Veblen TT (2001) Inter-hemispheric synchrony of forest fires and the El Niño–Southern Oscillation. *Global Ecology and Biogeography* **10**, 315–326. doi:10.1046/J.1466-822X.2001.00234.X
- Lenihan JL, Drapek R, Bachelet D, Neilson RP (2003) Climate change effects on vegetation distribution, carbon, and fire in California. *Ecological Applications* **13**, 1667–1681. doi:10.1890/025295
- Mann ME, Bradley RS, Hughes MK (1998) Global-scale temperature patterns and climate forcing over the past six centuries. *Nature* **392**, 779–787. doi:10.1038/33859
- Mann ME, Bradley RS, Hughes MK (2000) Long-term variability in the El Niño–Southern Oscillation and associated teleconnections. In ‘El Niño and the Southern Oscillation: multiscale variability and global and regional Impacts’. (Eds HF Diaz, V Markgraf) pp. 357–412. (Cambridge University Press: Cambridge)
- Mantua NJ, Hare SR, Zhang Y, Wallace JM, Francis RC (1997) A Pacific interdecadal climate oscillation with impacts on salmon production. *Bulletin of the American Meteorological Society* **78**, 1069–1079. doi:10.1175/1520-0477(1997)078<1069:APICOW>2.0.CO;2
- McCabe GJ, Dettinger MD (1999) Decadal variations in the strength of ENSO teleconnections with precipitation in the western United States. *International Journal of Climatology* **19**, 1399–1410. doi:10.1002/(SICI)1097-0088(19991115)19:13<1399::AID-JOC457>3.0.CO;2-A
- Miller RF, Heyerdahl EK, Fine-scale variation of historical fire regimes in sagebrush-steppe and juniper woodland: An example from California, USA. *International Journal of Wildland Fire* **17**, in press.
- Moody TJ, Fites-Kaufman J, Stephens SL (2006) Fire history and climate influences from forests in the northern Sierra Nevada, USA. *Fire Ecology* **2**, 115–141.
- Mooney CZ, Duval RD (1993) ‘Bootstrapping: a Non-Parametric Approach to Statistical Inference.’ (Sage Publications: Newbury Park, CA)
- Nagel TA, Taylor AH (2005) Fire and persistence of montane chaparral in mixed conifer forest landscapes in the northern Sierra Nevada, Lake Tahoe Basin, California, USA. *The Journal of the Torrey Botanical Society* **132**, 442–457. doi:10.3159/1095-5674(2005)132[442:FAPOMC]2.0.CO;2
- Nigam S, Barlow M, Berbery EH (1999) Analysis links Pacific decadal variability to drought and streamflow in the United States. *EOS, Transactions, American Geophysical Union* **80**, 621–625. doi:10.1029/99EO00412
- Norman SP, Taylor AH (2003) Tropical and north Pacific teleconnections influence fire regimes in pine-dominated forests of north-eastern California. *Journal of Biogeography* **30**, 1081–1092.
- Norman SP, Taylor AH (2005) Pine forest expansion along a forest-meadow ecotone in north-eastern California, USA. *Forest Ecology and Management* **215**, 51–68.
- Palmer WC (1965) Meteorological drought. US Department of Commerce, Weather Bureau Research Paper No. 45. (Washington, DC)
- Parker AJ (1991) Forest/environment relationships in Lassen Volcanic National Park, California, USA. *Journal of Biogeography* **18**, 534–552.
- Ritchie MW (2005) Ecological research at the Goosenest Adaptive Management Area in north-eastern California. USDA Forest Service, Pacific Southwest Research Station General Technical Report PSW-GTR-192. (Albany, CA)
- Sakulich J, Taylor AH (2007) Fire regimes and forest structure in a sky island mixed conifer forest, Guadalupe Mountains National Park, Texas, USA. *Forest Ecology and Management* **241**, 62–73. doi:10.1016/J.FORECO.2006.12.029
- Schoennagel T, Veblen TT, Romme WH, Sibold JS, Cook ER (2005) ENSO and PDO variability affect drought-induced fire occurrence in Rocky Mountain subalpine forests. *Ecological Applications* **15**, 2000–2014. doi:10.1890/04-1579
- Schonher T, Nicholson SE (1989) The relationship between California rainfall and ENSO events. *Journal of Climate* **2**(11), 1258–1269. doi:10.1175/1520-0442(1989)002<1258:TRBCRA>2.0.CO;2
- Schulz P (1954) ‘Indians of Lassen Volcanic National Park.’ (Loomis Museum Associates: Red Bluff, CA)
- Sibold JS, Veblen TT (2006) Relationships of subalpine forest fires in the Colorado Front Range with interannual and multidecadal-scale climatic variation. *Journal of Biogeography* **33**, 833–842. doi:10.1111/J.1365-2699.2006.01456.X
- Skinner CN, Chang C (1996) Fire regimes, past and present. In ‘Sierra Nevada Ecosystem Project: Final Report to Congress, Vol. II: Assessments and Scientific Basis for Management Options’. pp. 1041–1069. Water Resources Center Report No. 37. (Centers for Water and Wildland Resources, University of California: Davis, CA)
- Skinner CN, Taylor AH (2006) Southern Cascade bioregion. In ‘Fire in California’s ecosystems’. (Eds NS Sugihara, JW van Wagendonk, KE Shaffer, J Fites-Kaufman, AE Thode) pp. 195–224. (University of California Press: Berkeley, CA)
- Stephens SL, Skinner CN, Gill SJ (2003) A dendrochronology-based fire history of Jeffrey pine–mixed conifer forests in the Sierra San Pedro Martir, Mexico. *Canadian Journal of Forest Research* **33**, 1090–1101. doi:10.1139/X03-031
- Stokes MA, Smiley TL (1968) ‘An introduction to tree-ring dating.’ (University of Chicago Press: Chicago, IL)
- Strong DH (1973) ‘These happy hunting grounds, a history of the Lassen Region.’ (Walker Lithograph Inc.: Red Bluff, CA)
- Swetnam TW (1993) Fire history and climate change in giant sequoia groves. *Science* **262**, 885–889. doi:10.1126/SCIENCE.262.5135.885
- Swetnam TW, Baisan CH (2003) Tree-ring reconstructions of fire and climate history in the Sierra Nevada and south-western United States. In ‘Fire and climatic change in temperate ecosystems of the Western Americas’. (Eds TT Veblen, WL Baker, B Montenegro, TW Swetnam) pp. 158–195. (Springer Verlag: New York)
- Swetnam TW, Betancourt JL (1990) Fire–Southern Oscillation relations in the south-western United States. *Science* **249**, 1017–1020. doi:10.1126/SCIENCE.249.4972.1017
- Swetnam TW, Betancourt JL (1998) Mesoscale disturbance and ecological response to decadal climatic variability in the American South-west. *Journal of Climate* **11**, 3128–3147. doi:10.1175/1520-0442(1998)011<3128:MDAERT>2.0.CO;2
- Taylor AH (1990) Habitat segregation and regeneration patterns of red fir and mountain hemlock in ecotonal forests, Lassen Volcanic National Park, California. *Physical Geography* **11**, 36–48.
- Taylor AH (1995) Forest expansion and climate change in the mountain hemlock (*Tsuga mertensiana*) zone, Lassen Volcanic National Park, California, USA. *Arctic and Alpine Research* **27**, 207–216. doi:10.2307/1551951
- Taylor AH (2000) Fire regimes and forest changes in mid- and upper montane forests of the southern Cascades, Lassen Volcanic National Park, California, USA. *Journal of Biogeography* **27**, 87–104. doi:10.1046/J.1365-2699.2000.00353.X
- Taylor AH, Beaty RM (2005) Climatic influences on fire regimes in the northern Sierra Nevada mountains, Lake Tahoe Basin, Nevada, USA. *Journal of Biogeography* **32**, 425–438. doi:10.1111/J.1365-2699.2004.01208.X

- Taylor AH, Skinner CN (2003) Spatial patterns and controls on historical fire regimes and forest structure in the Klamath Mountains. *Ecological Applications* **13**, 704–719. doi:10.1890/1051-0761(2003)013[0704:SPACOH]2.0.CO;2
- Taylor AH, Solem MN (2001) Fire regimes and stand dynamics in an upper montane forest landscape in the southern Cascades, Caribou Wilderness, California. *Journal of the Torrey Botanical Society* **128**, 350–361. doi:10.2307/3088667
- Trouet V, Taylor AH, Carleton AM, Skinner CN (2006) Fire–climate interactions in forests of the American Pacific coast. *Geophysical Research Letters* **33**, L18704. doi:10.1029/2006GL027502
- Veblen TT, Kitzberger T, Donnegan J (2000) Climate and human influences on fire regimes in ponderosa pine forests in the Colorado Front Range. *Ecological Applications* **10**, 1178–1195. doi:10.1890/1051-0761(2000)010[1178:CAHIOF]2.0.CO;2
- Westerling A, Swetnam T (2003) Interannual to decadal drought and wildfire in the western United States. *EOS, Transactions, American Geophysical Union* **84**, 545–560. doi:10.1029/2003EO490001
- Westerling AL, Hidalgo HG, Cayan DR, Swetnam TW (2006) Warming and earlier spring increases western US forest wildfire activity. *Science* **313**, 940–943. doi:10.1126/SCIENCE.1128834
- Wilken G (1967) History and fire record of a timberland brushfield in the Sierra Nevada of California. *Ecology* **48**, 302–324. doi:10.2307/1933114

Manuscript received 19 February 2007, accepted 5 June 2007